

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
27 September 2001 (27.09.2001)

PCT

(10) International Publication Number  
**WO 01/70276 A2**

(51) International Patent Classification<sup>7</sup>: **A61K 48/00,**  
C12N 15/86

(US). ZABNER, Joseph; 712 McLean Street, Iowa City,  
IA 52246 (US).

(21) International Application Number: PCT/US01/09123

(74) Agents: SPRATT, Gwendolyn, D. et al.; Needle & Rosen-  
berg, P.C., 127 Peachtree Street, N.E., Suite 1200, Atlanta,  
GA 30303-1811 (US).

(22) International Filing Date: 22 March 2001 (22.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/533,427 22 March 2000 (22.03.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(71) Applicants: THE GOVERNMENT OF THE UNITED  
STATES OF AMERICA, as represented by THE  
SECRETARY, DEPARTMENT OF HEALTH AND  
HUMAN SERVICES [US/US]; C/o National Institutes  
of Health, Office of Technology Transfer, 6011 Executive  
Boulevard, Suite 325, Rockville, MD 20852-3804 (US).  
UNIVERSITY OF IOWA RESEARCH FOUNDA-  
TION [US/US]; 100 Oakdale Campus, #219 TIC, 219  
Technology Innovation Center, Iowa City, IA 52242 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished  
upon receipt of that report

(72) Inventors: CHIORINI, John, A.; 2604 Loma Street, Sil-  
ver Spring, MD 20902 (US). KOTIN, Robert, M.; 707  
Gormley, Rockville, MD 20850 (US). DAVIDSON, Bev-  
erly; 3640 Johnston Way, N.E., North Liberty, IA 52317

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.



WO 01/70276 A2

(54) Title: AAV5 VECTOR FOR TRANSDUCING BRAIN CELLS AND LUNG CELLS

(57) Abstract: The present invention provides methods of delivering nucleic acids to specific regions, tissues and cell types of the CNS. More particularly the invention provides methods of delivering nucleic acids to cells of the CNS such as cerebellar cells and ependymal cells. The invention also provides methods of delivering nucleic acids to cells of the lung such as alveolar cells using AAV5 vectors and particles.

## AAV5 VECTOR FOR TRANSDUCING BRAIN CELLS AND LUNG CELLS

This application claims priority from U.S. Application Serial No. 09/533,427,  
5 filed March 22, 2000 which is hereby incorporated in its entirety by this reference.

### BACKGROUND OF THE INVENTION

#### Field of the Invention

The present invention provides adeno-associated virus 5 (AAV5) and vectors  
10 derived therefrom. Thus, the present invention relates to AAV5 vectors for and  
methods of delivering nucleic acids to cells of subjects. More specifically, the present  
invention provides methods of delivering nucleic acids to cerebellar cells, ependymal  
cells, neurons astrocytes, airway epithelial cells and alveolar cells of subjects. More  
particularly, the invention provides methods of delivering nucleic acids to cerebellar  
15 cells, ependymal cells and alveolar cells.

#### Background Art

Adeno associated virus (AAV) is a small nonpathogenic virus of the  
parvoviridae family (for review see 28). AAV is distinct from the other members of  
20 this family by its dependence upon a helper virus for replication. In the absence of a  
helper virus, AAV has been shown to integrate in a locus specific manner into the q  
arm of chromosome 19 (21). The approximately 5 kb genome of AAV consists of one  
segment of single stranded DNA of either plus or minus polarity. Physically, the  
parvovirus virion is non-enveloped and its icosohedral capsid is approximately 20-25  
25 nm in diameter.

To date 8 serologically distinct AAVs have been identified and 6 have been  
isolated from humans or primates and are referred to as AAV types 1-6 (1). The most  
extensively studied of these isolates is AAV type 2 (AAV2). The genome of AAV2 is  
30 4680 nucleotides in length and contains two open reading frames (ORFs), the right

Rep52, Rep68 and Rep78, which are involved in regulation of replication and transcription in addition to the production of single-stranded progeny genomes (5-8, 11, 12, 15, 17, 19, 21-23, 25, 34, 37-40). Furthermore, two of the Rep proteins have been associated with the preferential integration of AAV genomes into a region of the q arm of human chromosome 19. Rep68/78 have also been shown to possess NTP binding activity as well as DNA and RNA helicase activities. The Rep proteins possess a nuclear localization signal as well as several potential phosphorylation sites. Mutation of one of these kinase sites resulted in a loss of replication activity.

10       The ends of the genome are short inverted terminal repeats which have the potential to fold into T-shaped hairpin structures that serve as the origin of viral DNA replication. Within the ITR region two elements have been described which are central to the function of the ITR, a GAGC repeat motif and the terminal resolution site (TRS). The repeat motif has been shown to bind Rep when the ITR is in either a linear or  
15       hairpin conformation (7, 8, 26).

          This binding serves to position Rep68/78 for cleavage at the TRS which occurs in a site- and strand-specific manner. In addition to their role in replication, these two elements appear to be central to viral integration. Contained within the chromosome 19  
20       integration locus is a Rep binding site with an adjacent TRS. These elements have been shown to be functional and necessary for locus specific integration.

          The AAV2 virion is a non-enveloped, icosohedral particle approximately 20-25 nm in diameter. The capsid is composed of three related proteins referred to as VP1,2  
25       and 3 which are encoded by the right ORF. These proteins are found in a ratio of 1:1:10 respectively. The capsid proteins differ from each other by the use of alternative splicing and an unusual start codon. Deletion analysis of has shown that removal or alteration of AAV2 VP1 which is translated from an alternatively spliced message results in a reduced yield of infectious particles (15, 16, 38). Mutations within the VP3  
30       coding region result in the failure to produce any single-stranded progeny DNA or infectious particles (15, 16, 38).

The following features of the characterized AAVs have made them attractive vectors for gene transfer (16). AAV vectors have been shown *in vitro* to stably integrate into the cellular genome; possess a broad host range; transduce both dividing and non dividing cells *in vitro* and *in vivo* (13, 20, 30, 32) and maintain high levels of expression of the transduced genes (41). Viral particles are heat stable, resistant to solvents, detergents, changes in pH, temperature, and can be concentrated on CsCl gradients (1,2). Integration of AAV provirus is not associated with any long term negative effects on cell growth or differentiation (3,42). The ITRs have been shown to be the only cis elements required for replication, packaging and integration (35) and may contain some promoter activities (14).

AAV2 was originally thought to infect primate and non-primate cell types provided the appropriate helper virus was present. However, the inability of AAV2 to infect certain cell types is now known to be due to the particular cellular tropism exhibited by the AAV2 virus. Recent work has shown that some cell lines are transduced very poorly by AAV2 (30). Binding studies have indicated that heparin sulfate proteoglycans are necessary for high efficiency transduction with AAV2. AAV5 is a unique member of the parvovirus family. The present DNA hybridization data indicate a low level of homology with the published AAV1-4 sequences (31). The present invention shows that, unlike AAV2, AAV5 transduction is not effected by heparin as AAV2 is and therefore will not be restricted to the same cell types as AAV2.

The present invention provides a vector comprising the AAV5 virus or a vector comprising subparts of the virus, as well as AAV5 viral particles. While AAV5 is similar to AAV2, the two viruses are found herein to be physically and genetically distinct. These differences endow AAV5 with some unique properties and advantages which better suit it as a vector for gene therapy. For example, one of the limiting features of using AAV2 as a vector for gene therapy is production of large amounts of virus. Using standard production techniques, AAV5 is produced at a 10-50 fold higher level compared to AAV2. Because of its unique TRS site and rep proteins, AAV5 should also have a distinct integration locus compared to AAV2.



Furthermore, as shown herein, AAV5 capsid protein, again surprisingly, is distinct from AAV2 capsid protein and exhibits different tissue tropism, thus making AAV5 capsid-containing particles suitable for transducing cell types for which AAV2 is unsuited or less well-suited. AAV2 and AAV5 have been shown to be serologically  
5 distinct and thus, in a gene therapy application, AAV5, and AAV5-derived vectors, would allow for transduction of a patient who already possess neutralizing antibodies to AAV2 either as a result of natural immunological defense or from prior exposure to AAV2 vectors. Another advantage of AAV5 is that AAV5 cannot be rescued by other serotypes. Only AAV5 can rescue the integrated AAV5 genome and effect replication,  
10 thus avoiding unintended replication of AAV5 caused by other AAV serotypes. Thus, the present invention, by providing these new recombinant vectors and particles based on AAV5 provides a new and highly useful series of vectors.

#### SUMMARY OF THE INVENTION

15

The present invention provides methods of delivering a nucleic acid to specific regions, tissues and cell types of the central nervous system (CNS) such as ependymal cells, cerebellar cells, neurons, and astrocytes. In particular, the nucleic acids are delivered to specific regions and cells of the brain, particularly, ependymal cells and  
20 cerebellar cells.

The present invention also provides methods of delivering a nucleic acid to lung cells such as alveolar cells.

25 The present invention provides a nucleic acid vector comprising a pair of adeno-associated virus 5 (AAV5) inverted terminal repeats and a promoter between the inverted terminal repeats.

The present invention further provides an AAV5 particle containing a vector  
30 comprising a pair of AAV2, AAV4 or AAV5 inverted terminal repeats.

Additionally, the instant invention provides an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:1 (AAV5 genome). Furthermore, the present invention provides an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:1 (AAV5 genome).

5

The present invention provides an isolated nucleic acid encoding an AAV5 Rep protein, for example, the nucleic acid as set forth in SEQ ID NO:10. Additionally provided is an isolated full-length AAV5 Rep protein or a unique fragment thereof. Additionally provided is an isolated AAV5 Rep 40 protein having the amino acid  
10 sequence set forth in SEQ ID NO:12, or a unique fragment thereof. Additionally provided is an isolated AAV5 Rep 52 protein having the amino acid sequence set forth in SEQ ID NO:2, or a unique fragment thereof. Additionally provided is an isolated AAV5 Rep 68 protein, having the amino acid sequence set forth in SEQ ID NO:14 or a unique fragment thereof. Additionally provided is an isolated AAV5 Rep 78 protein  
15 having the amino acid sequence set forth in SEQ ID NO:3, or a unique fragment thereof. The sequences for these proteins are provided below in the Sequence Listing and elsewhere in the application where the proteins are described.

The present invention further provides an isolated AAV5 capsid protein, VP1,  
20 having the amino acid sequence set forth in SEQ ID NO:4, or a unique fragment thereof. Additionally provided is an isolated AAV5 capsid protein, VP2, having the amino acid sequence set forth in SEQ ID NO:5, or a unique fragment thereof. Also provided is an isolated AAV5 capsid protein, VP3, having the amino acid sequence set forth in SEQ ID NO:6, or a unique fragment thereof.

25

The present invention additionally provides an isolated nucleic acid encoding AAV5 capsid protein, for example, the nucleic acid set forth in SEQ ID NO:7, or a unique fragment thereof.

The present invention provides an isolated nucleic acid encoding an AAV5 inverted terminal repeat, for example, the nucleic acid set forth in SEQ ID NO: 19 or SEQ ID NO: 20, or a unique fragment thereof.

5           The present invention further provides an AAV5 particle comprising a capsid protein consisting essentially of the amino acid sequence set forth in SEQ ID NO:4, or a unique fragment thereof.

            Additionally provided by the present invention is an isolated nucleic acid  
10   comprising an AAV5 p5 promoter having the nucleic acid sequence set forth in SEQ ID NO:18, or a unique fragment thereof.

            The instant invention provides a method of screening a cell for infectivity by AAV5 comprising contacting the cell with AAV5 and detecting the presence of AAV5  
15   in the cells.

            The present invention further provides a method of delivering a nucleic acid to a cell comprising administering to the cell an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats,  
20   thereby delivering the nucleic acid to the cell.

            The present invention also provides a method of delivering a nucleic acid to a subject comprising administering to a cell from the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats,  
25   and returning the cell to the subject, thereby delivering the nucleic acid to the subject.

            The present invention also provides a method of delivering a nucleic acid to a cell in a subject comprising administering to the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby  
30   delivering the nucleic acid to a cell in the subject.

The instant invention further provides a method of delivering a nucleic acid to a cell in a subject having antibodies to AAV2 comprising administering to the subject an AAV5 particle comprising the nucleic acid, thereby delivering the nucleic acid to a cell in the subject.

5

### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows Heparin inhibition results. Cos cells were plated in 12 well dishes at  $5 \times 10^4$  cells per well. Serial dilutions of AAV2 or AAV5 produced and  
10 purified as previously described and supplemented with  $5 \times 10^5$  particles of wt adenovirus were incubated for 1 hr at Rt in the presence of 20  $\mu\text{g/ml}$  heparin (sigma). Following this incubation the virus was added to the cells in 400  $\mu\text{l}$  of media for 1 hr after which the media was removed, the cells rinsed and fresh media added. After 24 hrs the plates were stained for Bgal activity.

15

Figure 2 shows AAV2 and AAV5 vector and helper complementation. Recombinant AAV particles were produced as previously described using a variety of vector and helper plasmids as indicated the bottom of the graph. The vector plasmids contained the Bgal gene with and RSV promoter and flanked by either AAV2 ITRs  
20 (2ITR) or AAV5 ITRs (5ITR). The helper plasmids tested contained either AAV2 Rep and cap genes (2repcap) AAV5 rep and cap genes with or without an SV40 promoter (5repcapA and 5repcapb respectively) only the AAV2 rep gene (2rep) in varying amounts (1) or (.5) or an empty vector (pUC). The resulting AAV particles were then titrated on cos cells. AAV particles were only produced when the same serotype of ITR  
25 and Rep were present.

Figure 3 shows AAV2 and AAV5 tissue tropism. Transduction of a variety of cell types indicated that AAV2 and AAV5 transduce cells with different efficiencies. Equal number of either AAV2 or AAV5 particles were used to transduce a variety of  
30 cell types and the number of bgal positive cells is reported.

Figure 4 is a sequence comparison of the AAV2 genome and the AAV5 genome.

Figure 5 is a sequence comparison of the AAV2 VP1 capsid protein and the  
5 AAV5 VP1 capsid protein.

Figure 6 is a sequence comparison of the AAV2 rep 78 protein and the AAV5 rep 78 protein.

10 Figure 7 shows the transduction of airway epithelial cells by AAV5. Primary airway epithelial cells were cultured and plated. Cells were transduced with an equivalent number of rAAV2 or rAAV5 particles containing a nuclear localized  $\beta$ -gal transgene with 50 particles of virus/cell (MOI 50) and continued in culture for 10 days.  $\beta$ -gal activity was determined and the relative transduction efficiency compared.  
15 AAV5 transduced these cells 50- fold more efficiently than AAV2. This is the first time apical cells or cells exposed to the air have been shown to be infected by a gene therapy agent.

Figure 8 shows transduction of striated muscle by AAV5. Chicken myoblasts  
20 were cultured and plated. Cells were allowed to fuse and then transduced with a similar number of particles of rAAV2 or rAAV5 containing a nuclear localized  $\beta$ -gal transgene after 5 days in culture. The cells were stained for  $\beta$ -gal activity and the relative transduction efficiency compared. AAV5 transduced these cells approximately 16 fold more efficiently than AAV2.

25 Figure 9 shows transduction of rat brain explants by AAV5. Primary neonatal rat brain explants were prepared. After 7 days in culture, cells were transduced with a similar number of particles of rAAV5 containing a nuclear localized  $\beta$ -gal transgene. After 5 days in culture, the cells were stained for  $\beta$ -gal activity. Transduction was  
30 detected in a variety of cell types including astrocytes, neuronal cells and glial cells.

Figure 10 shows transduction of human umbilical vein endothelial cells by AAV5. Human umbilical vein endothelial cells were cultured and plated. Cells were transduced with rAAV2 or rAAV5 containing a nuclear localized  $\beta$ -gal transgene with 10 particles of virus/ cell (MOI 5) in minimal media then returned to complete media. After 24 hrs in culture, the cells were stained for  $\beta$ -gal activity and the relative transduction efficiency compared. As shown in AAV5 transduced these cell 5-10 fold more efficiently than AAV2.

Figure 11 shows the number of  $\beta$ -galactosidase positive cells after cerebral injection. Animals were injected with rAAV2 $\beta$ gal, rAAV4 $\beta$ gal, or rAAV5 $\beta$ gal into the ventricle or striatum of mice, and brains taken at the times indicated. Blocks (2.6 mm, surrounding the injection site) were sectioned, processed for  $\beta$ -galactosidase histochemistry, and transgene positive cells counted. Data represent mean  $\pm$  SEM. \*,  $p < 0.05$ , \*\* $p < 0.005$

Figure 12 shows The distribution of  $\beta$ -galactosidase positive cells in brains of mice at 3 or 15 weeks following injection of rAAV2 $\beta$ gal, rAAV4 $\beta$ gal or rAAV5 $\beta$ gal.  $\beta$ -galactosidase positive cells within the ependymal, striatal, or 'other' (septal and fornix regions, corpus callosum, and neocortex) regions, from sections encompassing the injection site, were counted and that value represented as a percent of the total number of transgene positive cells. (A) Data from sections obtained 3 or 15 weeks after intraventricular injection. (B) Data from sections obtained 3 or 15 weeks after injection into the striatum. Data represent mean  $\pm$  SEM.

Figure 13 shows the  $\beta$ -galactosidase histochemistry for transgene positive cells after striatal injection of rAAV vectors. (A,B) Representative photomicrograph of sections from mice injected with rAAV2 $\beta$ gal. Panel B is a magnified photograph of the transgene positive striatal region seen in A. (C) Demonstration of distinctive ependymal-specific staining for  $\beta$ -galactosidase in sections from animals injected with rAAV4 $\beta$ gal. (D-F). Photomicrographs illustrating the extensive distribution of transgene positive cells after rAAV5 $\beta$ gal injection. (E) Magnification of the striatal

region in D. (F) Transgene positive cells in the cortex distant from the injection site. The photomicrographs are representative of at least three independent experiments. fi, fimbria region; lv, lateral ventricle; sp, medial septal region; st, striatum.

5           Figure 14 shows the identification of transduced cells after intrastriatal injection of rAAV5 $\beta$ gal. Fifteen weeks after injection of rAAV5 $\beta$ gal coronal brain sections were dual stained for  $\beta$ -galactosidase (green nuclei) and NeuN (neuronal specific, red nuclei and light red cytoplasm), or for  $\beta$ -galactosidase and GFAP (astrocyte-specific, red cell processes). Confocal microscopy image analysis was performed and representative  
10 two-color merged images of single z-series slices are shown. In the striatum, both transduced neurons (yellow cell nuclei in A) and transduced astrocytes (B) were detected. In the medial septal region transduction appeared to be restricted to neurons (C), while in the corpus callosum the transduced cells were GFAP positive astrocytes (D). Images were captured using a 40X (A,B,D) or 63X (C) oil immersion objective.

15           Figure 15 shows gene transfer to the apical surface of well-differentiated human airway epithelia by different recombinant AAV serotypes. (A,D) *Enface* images of human airway epithelia (A) and epithelia transduced with 500 particles per cell of AAV2/ $\beta$ Gal (B), AAV4/ $\beta$ Gal (C) and AAV5/ $\beta$ Gal (D). The blue staining show cells  
20 that have been transduced with vector. Fig 15E shows the quantitative  $\beta$ -gal activity of airway epithelia infected the different recombinant AAV serotypes (AAV2/ $\beta$ Gal, AAV4/ $\beta$ Gal, and AAV5/ $\beta$ Gal). Data are the mean  $\beta$ -gal activity per mg protein  $\pm$  SEM (n = 4-12). Asterisk indicates p < 0.01.

25           Figure 16 shows the binding of AAV2/ $\beta$ Gal, AAV4/ $\beta$ Gal, and AAV5/ $\beta$ Gal to organotypic cultures of ciliated human airway epithelia. A. Figure shows the dot blot of virus bound to the epithelia of 3 experiments with seven epithelia per experiment (input virus 500 particles/cell). For the purpose of quantitation, a dilution series of rAAV plasmid was also blotted and probed to demonstrate the linearity of the detection  
30 system. B. Figure shows the results of the quantification of the dot blot data. The data

are means  $\pm$  SEM of the percentage of total virus added that remained epithelia-associated after a 30 min incubation. Asterisk indicates  $p < 0.05$ .

Figure 17 shows the effect of dose on AAV2/ $\beta$ Gal and AAV5/ $\beta$ Gal-mediated  
5 gene transfer to human airway epithelia. Human airway epithelia were exposed to increasing number of particles per cell of AAV2/ $\beta$ Gal and AAV5/ $\beta$ Gal from the apical surface. The epithelia were then rinsed after 60 min and incubated for 2 weeks prior to analysis of  $\beta$ -galactosidase activity. Data are the  $\beta$ -gal activity per mg protein  $\pm$  SEM (n = 4).

10

Figure 18 shows the effect of incubation time on AAV5/ $\beta$ Gal-mediated gene transfer to human airway epithelia. Human airway epithelia were exposed to 500 particles per cell of AAV5/ $\beta$ Gal from the apical surface. The epithelia were then rinsed after 30, 60 min or 90, 360 and 720 min and incubated for 2 weeks prior to analysis of  
15  $\beta$ -galactosidase activity. Data are the  $\beta$ -gal activity per mg protein  $\pm$  SEM (n = 4). Asterisk indicates  $p < 0.01$ .

Figure 19A,B shows the effect of soluble heparin on AAV gene transfer to human airway epithelia. To compete off AAV binding and gene transfer, in some  
20 conditions the viruses were pretreated with 20  $\mu$ g/ml of soluble heparin for 30 min. Fig 19A shows the effect of heparin on AAV gene transfer to human airway epithelia from the apical side and Fig 5B from the basolateral side. Five hundred particles per cell of either AAV2/ $\beta$ Gal or AAV5/ $\beta$ Gal were added for 30 min at 4°C.  $\beta$ -galactosidase was measured 14 days later. Data are mean  $\pm$  SEM, n=8 in each group. Asterisk in Fig 19B  
25 indicates  $p = 0.018$ .

Figure 20 shows the AAV5/ $\beta$ Gal-mediated gene transfer to murine conducting airway epithelia, and alveolar epithelia *in vivo*. Mice were exposed to  $1 \times 10^{10}$  particles of either AAV2/ $\beta$ Gal or AAV5/ $\beta$ Gal via nasal instillation. After 30 days the mice were  
30 sacrificed, the lungs were fixed and stained with X-Gal. Fig. 20 A,B shows representative photomicrographs showing ciliated and non-ciliated cells transduced by



AAV2/ $\beta$ Gal (A) and AAV5/ $\beta$ Gal (B). Figure 20 C shows quantitation of gene transfer by number of blue nuclei of  $\beta$ gal-expressing bronchial and alveolar cells per microscopic field.  $n = 5$  mice per group. Asterisk indicates  $p < 0.01$ .

5        Figure 21 shows a sagittal section of a mouse cerebellum injected with AAV5 expressing nuclear targeted  $\beta$ -galactosidase driven off an RSV promoter. At 7 weeks postinjection, the animal was deeply anesthetized and s transcardially perfused with 4% paraformaldehyde. Cerebellum was sectioned at 50  $\mu$ m thickness and sections were processed for X-gal histochemistry. AAV5 transduced large numbers of Purkinje cells,  
10    stellate and basket neurons and a smaller number of Golgi neurons.

Figure 22 -double label immunofluorescence showed that within the cerebellar cortex AAV5 transduced neurons but not glia. This section is from the same animal as A and processed with antibodies against glial fibrillary acid protein (GFAP) which is  
15    red and  $\beta$ -galactosidase which fluoresces green. Arrow points to a typical  $\beta$ -galactosidase positive cell. Confocal microscopy shows that there is no colocalization between GFAP and AAV5  $\beta$ -galactosidase. Thus all cells which are transduced are nonglial (neurons).

20        Figure 23 shows a third cerebellar section from the same animal as Fig. 21 and Fig. 22 which was processed for double label immunofluorescence with antibodies against calbindin (red) and  $\beta$ -galactosidase (green). Calbindin is expressed in Purkinje cells but not other cerebellar neurons. Confocal microscopy showed strong colocalization between calbindin and the AAV5 expressed  $\beta$ -galactosidase in the  
25    Purkinje cell monolayer (arrow). However there were many  $\beta$ -galactosidase positive neurons in molecular layer and granule cell layer (example arrowhead) that did not express calbindin, confirming that several classes of neurons had been transduced.

## DETAILED DESCRIPTION OF THE INVENTION

30

As used in the specification and in the claims, "a" can mean one or more,

depending upon the context in which it is used. The terms “having” and “comprising” are used interchangeably herein, and signify open ended meaning.

The present application provides a recombinant adeno-associated virus 5 (AAV5). This virus has one or more of the characteristics described below. The compositions of the present invention do not include wild-type AAV5. The methods of the present invention can use either wild-type AAV5 or recombinant AAV5-based delivery.

10 The present invention provides novel AAV5 particles, recombinant AAV5 vectors, recombinant AAV5 virions and novel AAV5 nucleic acids and polypeptides.

An AAV5 particle is a viral particle comprising an AAV5 capsid protein. A recombinant AAV5 vector is a nucleic acid construct that comprises at least one unique nucleic acid of AAV5. A recombinant AAV5 virion is a particle containing a  
15 recombinant AAV5 vector, wherein the particle can be either an AAV5 particle as described herein or a non-AAV5 particle. Alternatively, the recombinant AAV5 virion is an AAV5 particle containing a recombinant vector, wherein the vector can be either an AAV5 vector as described herein or a non-AAV5 vector. These vectors, particles, virions, nucleic acids and polypeptides are described below.

20

The present invention provides the nucleotide sequence of the AAV5 genome and vectors and particles derived therefrom. Specifically, the present invention provides a nucleic acid vector comprising a pair of AAV5 inverted terminal repeats (ITRs) and a promoter between the inverted terminal repeats. While the rep proteins of  
25 AAV2 and AAV5 will bind to either a type 2 ITR or a type 5 ITR, efficient genome replication only occurs when type 2 Rep replicates a type 2 ITR and a type 5 Rep replicates a type 5 ITR. This specificity is the result of a difference in DNA cleavage specificity of the two Reps which is necessary for replication. AAV5 Rep cleaves at CGGT<sup>^</sup>GTGA (SEQ ID NO: 21) and AAV2 Rep cleaves at CGGT<sup>^</sup>TGAG (SEQ ID  
30 NO: 22) (Chiorini et al., 1999. J. Virol. 73 (5) 4293-4298). Mapping of the AAV5 ITR terminal resolution site (TRS) identified this distinct cleavage site, CGGT<sup>^</sup>GTGA,

which is absent from the ITRs of other AAV serotypes. Therefore, the minimum sequence necessary to distinguish AAV5 from AAV2 is the TRS site where Rep cleaves in order to replicate the virus. Examples of the type 5 ITRs are shown in SEQ ID NO: 19 and SEQ ID NO: 20, AAV5 ITR "flip" and AAV5 "flop", respectively.

- 5 Minor modifications in an ITR of either orientation are contemplated and are those that will not interfere with the hairpin structure formed by the AAV5 ITR as described herein. Furthermore, to be considered within the term "AAV5 ITR" the nucleotide sequence must retain one or more features described herein that distinguish the AAV5 ITR from the ITRs of other serotypes, e.g. it must retain the Rep binding site described  
10 herein.

- The D- region of the AAV5 ITR (SEQ ID NO: 23), a single stranded region of the ITR, inboard of the TRS site, has been shown to bind a factor which depending on  
15 its phosphorylation state correlates with the conversion of the AAV from a single stranded genome to a transcriptionally active form that allows for expression of the viral DNA. This region is conserved between AAV2, 3, 4, and 6 but is divergent in AAV5. The D+ region is the reverse complement of the D- region.

- 20 The promoter can be any desired promoter, selected by known considerations, such as the level of expression of a nucleic acid functionally linked to the promoter and the cell type in which the vector is to be used. That is, the promoter can be tissue/cell-specific. Promoters can be prokaryotic, eukaryotic, fungal, nuclear, mitochondrial, viral or plant promoters. Promoters can be exogenous or endogenous to the cell type  
25 being transduced by the vector. Promoters can include, for example, bacterial promoters, known strong promoters such as SV40 or the inducible metallothionein promoter, or an AAV promoter, such as an AAV p5 promoter. Additionally, chimeric regulatory promoters for targeted gene expression can be utilized. Examples of these regulatory systems, which are known in the art, include the tetracycline based  
30 regulatory system which utilizes the tet transactivator protein (tTA), a chimeric protein containing the VP16 activation domain fused to the tet repressor of *Escherichia coli*,

the IPTG based regulatory system, the CID based regulatory system, and the Ecdysone based regulatory system (44). Other promoters include promoters derived from actin genes, immunoglobulin genes, cytomegalovirus (CMV), adenovirus, bovine papilloma virus, adenoviral promoters, such as the adenoviral major late promoter, an inducible  
5 heat shock promoter, respiratory syncytial virus, Rous sarcomas virus (RSV), etc., specifically, the promoter can be AAV2 p5 promoter or AAV5 p5 promoter. More specifically, the AAV5 p5 promoter can be about same location in SEQ ID NO: 1 as the AAV2 p5 promoter, in the corresponding AAV2 published sequence. An example of an AAV5 p5 promoter is nucleotides 220-338 of SEQ ID NO: 1. Additionally, the  
10 p5 promoter may be enhanced by nucleotides 1-130 of SEQ ID NO:1. Furthermore, smaller fragments of p5 promoter that retain promoter activity can readily be determined by standard procedures including, for example, constructing a series of deletions in the p5 promoter, linking the deletion to a reporter gene, and determining whether the reporter gene is expressed, *i.e.*, transcribed and/or translated. The promoter  
15 can be the promoter of any of the AAV serotypes, and can be the p19 promoter (SEQ ID NO: 16) or the p40 promoter set forth in the sequence listing as SEQ ID NO: 17.

It should be recognized that any errors in any of the nucleotide sequences disclosed herein can be corrected, for example, by using the hybridization procedure  
20 described below with various probes derived from the described sequences such that the coding sequence can be reisolated and resequenced. Rapid screening for point mutations can also be achieved with the use of polymerase chain reaction-single strand conformation polymorphism (PCR-SSCP) (43). The corresponding amino acid sequence can then be corrected accordingly.

25

The AAV5-derived vector of the invention can further comprise a heterologous nucleic acid functionally linked to the promoter. By "heterologous nucleic acid" is meant that any heterologous or exogenous nucleic acid, *i.e.* not normally found in wild-type AAV5 can be inserted into the vector for transfer into a cell, tissue or organism.  
30 By "functionally linked" is meant that the promoter can promote expression of the heterologous nucleic acid, as is known in the art, and can include the appropriate

orientation of the promoter relative to the heterologous nucleic acid. Furthermore, the heterologous nucleic acid preferably has all appropriate sequences for expression of the nucleic acid. The nucleic acid can include, for example, expression control sequences, such as an enhancer, and necessary information processing sites, such as ribosome  
5 binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences.

The heterologous nucleic acid can encode beneficial proteins or polypeptides that replace missing or defective proteins required by the cell or subject into which the  
10 vector is transferred or can encode a cytotoxic polypeptide that can be directed, *e.g.*, to cancer cells or other cells whose death would be beneficial to the subject. The heterologous nucleic acid can also encode antisense RNAs that can bind to, and thereby inactivate, mRNAs made by the subject that encode harmful proteins. The heterologous nucleic acid can also encode ribozymes that can effect the  
15 sequence-specific inhibition of gene expression by the cleavage of mRNAs. In one embodiment, antisense polynucleotides can be produced from a heterologous expression cassette in an AAV5 vector construct where the expression cassette contains a sequence that promotes cell-type specific expression (Wirak *et al.*, *EMBO* 10:289 (1991)). For general methods relating to antisense polynucleotides, see *Antisense RNA and DNA*, D. A. Melton, Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY  
20 (1988).

Examples of heterologous nucleic acids which can be administered to a cell or subject as part of the present AAV5 vector can include, but are not limited to the  
25 following: nucleic acids encoding secretory and nonsecretory proteins, nucleic acids encoding therapeutic agents, such as tumor necrosis factors (TNF), such as TNF- $\alpha$ ; interferons, such as interferon- $\alpha$ , interferon- $\beta$ , and interferon- $\gamma$ ; interleukins, such as IL-1, IL-1 $\beta$ , and ILs -2 through -14; GM-CSF; adenosine deaminase; cellular growth factors, such as lymphokines; soluble CD4; Factor VIII; Factor IX; T-cell receptors;  
30 LDL receptor; ApoE; ApoC; alpha-1 antitrypsin; ornithine transcarbamylase (OTC); cystic fibrosis transmembrane receptor (CFTR); insulin; anti-apoptotic gene products;

proteins promoting neuronal survival, such as growth factors and glutamate receptors; Fc receptors for antigen binding domains of antibodies, such as immunoglobulins; anti-HIV decoy tar elements; and antisense sequences which inhibit viral replication, such as antisense sequences which inhibit replication of hepatitis B or hepatitis non-A, non-B virus. The nucleic acid is chosen considering several factors, including the cell to be transfected. Where the target cell is a blood cell, for example, particularly useful nucleic acids to use are those which allow the blood cells to exert a therapeutic effect, such as a gene encoding a clotting factor for use in treatment of hemophilia. Another target cell is the lung airway cell, which can be used to administer nucleic acids, such as those coding for the cystic fibrosis transmembrane receptor, which could provide a gene therapeutic treatment for cystic fibrosis. Other target cells include muscle cells where useful nucleic acids, such as those encoding cytokines and growth factors, can be transduced and the protein the nucleic acid encodes can be expressed and secreted to exert its effects on other cells, tissues and organs, such as the liver. Furthermore, the nucleic acid can encode more than one gene product, limited only, if the nucleic acid is to be packaged in a capsid, by the size of nucleic acid that can be packaged.

Furthermore, suitable nucleic acids can include those that, when transferred into a primary cell, such as a blood cell, cause the transferred cell to target a site in the body where that cell's presence would be beneficial. For example, blood cells such as TIL cells can be modified, such as by transfer into the cell of a Fab portion of a monoclonal antibody, to recognize a selected antigen. Another example would be to introduce a nucleic acid that would target a therapeutic blood cell to tumor cells. Nucleic acids useful in treating cancer cells include those encoding chemotactic factors which cause an inflammatory response at a specific site, thereby having a therapeutic effect.

Cells, particularly blood cells, muscle cells, airway epithelial cells, brain cells and endothelial cells having such nucleic acids transferred into them can be useful in a variety of diseases, syndromes and conditions. For example, suitable nucleic acids include nucleic acids encoding soluble CD4, used in the treatment of AIDS and  $\alpha$ -antitrypsin, used in the treatment of emphysema caused by  $\alpha$ -antitrypsin deficiency.

Other diseases, syndromes and conditions in which such cells can be useful include, for example, adenosine deaminase deficiency, sickle cell deficiency, brain disorders such as Alzheimer's disease, Huntington's disease, lysosomal storage diseases, Gaucher's disease, Hurler's disease, Krabbe's disease, motor neuron diseases such as

5    amyotrophic lateral sclerosis and dominant spinal cerebellar ataxias (examples include SCA1, SCA2, and SCA3), thalassemia, hemophilia, diabetes, phenylketonuria, growth disorders and heart diseases, such as those caused by alterations in cholesterol metabolism, and defects of the immune system.

10            As another example, hepatocytes can be transfected with the present vectors having useful nucleic acids to treat liver disease. For example, a nucleic acid encoding OTC can be used to transfect hepatocytes (*ex vivo* and returned to the liver or *in vivo*) to treat congenital hyperammonemia, caused by an inherited deficiency in OTC. Another example is to use a nucleic acid encoding LDL to target hepatocytes *ex vivo* or *in vivo*

15    to treat inherited LDL receptor deficiency. Such transfected hepatocytes can also be used to treat acquired infectious diseases, such as diseases resulting from a viral infection. For example, transduced hepatocyte precursors can be used to treat viral hepatitis, such as hepatitis B and non-A, non-B hepatitis, for example by transducing the hepatocyte precursor with a nucleic acid encoding an antisense RNA that inhibits

20    viral replication. Another example includes transferring a vector of the present invention having a nucleic acid encoding a protein, such as  $\alpha$ -interferon, which can confer resistance to the hepatitis virus.

For a procedure using transfected hepatocytes or hepatocyte precursors,

25    hepatocyte precursors having a vector of the present invention transferred in can be grown in tissue culture, removed from the tissue culture vessel, and introduced to the body, such as by a surgical method. In this example, the tissue would be placed directly into the liver, or into the body cavity in proximity to the liver, as in a transplant or graft. Alternatively, the cells can simply be directly injected into the liver, into the portal

30    circulatory system, or into the spleen, from which the cells can be transported to the liver via the circulatory system. Furthermore, the cells can be attached to a support,

such as microcarrier beads, which can then be introduced, such as by injection, into the peritoneal cavity. Once the cells are in the liver, by whatever means, the cells can then express the nucleic acid and/or differentiate into mature hepatocytes which can express the nucleic acid.

5

The AAV5-derived vector can include any normally occurring AAV5 sequences in addition to an ITR and promoter. Examples of vector constructs are provided below.

The present vector or AAV5 particle or recombinant AAV5 virion can utilize  
10 any unique fragment of the present AAV5 nucleic acids, including the AAV5 nucleic acids set forth in SEQ ID NOS: 1 and 7-11, 13, 15, 16, 17, and 18. To be unique, the fragment must be of sufficient size to distinguish it from other known sequences, most readily determined by comparing any nucleic acid fragment to the nucleotide sequences of nucleic acids in computer databases, such as GenBank. Such comparative searches  
15 are standard in the art. Typically, a unique fragment useful as a primer or probe will be at least about 8 or 10, preferable at least 20 or 25 nucleotides in length, depending upon the specific nucleotide content of the sequence. Additionally, fragments can be, for example, at least about 30, 40, 50, 75, 100, 200 or 500 nucleotides in length and can encode polypeptides or be probes. The nucleic acid can be single or double stranded,  
20 depending upon the purpose for which it is intended. Where desired, the nucleic acid can be RNA.

The present invention further provides an isolated AAV5 capsid protein to contain the vector. In particular, the present invention provides not only a polypeptide  
25 comprising all three AAV5 coat proteins, *i.e.*, VP1, VP2 and VP3, but also a polypeptide comprising each AAV5 coat protein individually, SEQ ID NOS: 4, 5, and 6, respectively. Thus an AAV5 particle comprising an AAV5 capsid protein comprises at least one AAV5 coat protein VP1, VP2 or VP3. An AAV5 particle comprising an AAV5 capsid protein can be utilized to deliver a nucleic acid vector to a cell, tissue or  
30 subject. For example, the herein described AAV5 vectors can be encapsidated in an AAV5 capsid-derived particle and utilized in a gene delivery method. Furthermore,



other viral nucleic acids can be encapsidated in the AAV5 particle and utilized in such delivery methods. For example, an AAV1, 2,3,4, or 6 vector (e.g. AAV1,2,3,4, or 6 ITR and nucleic acid of interest) can be encapsidated in an AAV5 particle and administered. Furthermore, an AAV5 chimeric capsid incorporating both AAV2 capsid and AAV5  
5 capsid sequences can be generated, by standard cloning methods, selecting regions from the known sequences of each protein as desired. For example, particularly antigenic regions of the AAV2 capsid protein can be replaced with the corresponding region of the AAV5 capsid protein. In addition to chimeric capsids incorporating AAV2 capsid sequences, chimeric capsids incorporating AAV1, 3, 4, or 6 and AAV5 capsid  
10 sequences can be generated, by standard cloning methods, selecting regions from the known sequences of each protein as desired.

The capsids can also be modified to alter their specific tropism by genetically altering the capsid to encode a specific ligand to a cell surface receptor. Alternatively,  
15 the capsid can be chemically modified by conjugating a ligand to a cell surface receptor. By genetically or chemically altering the capsids, the tropism can be modified to direct AAV5 to a particular cell or population of cells. The capsids can also be altered immunologically by conjugating the capsid to an antibody that recognizes a specific protein on the target cell or population of cells.

20

The capsids can also be assembled into empty particles by expression in mammalian, bacterial, fungal or insect cells. For example, AAV2 particles are known to be made from VP3 and VP2 capsid proteins in baculovirus. The same basic protocol can produce an empty AAV5 particle comprising an AAV5 capsid protein.

25

The herein described recombinant AAV5 nucleic acid derived vector can be encapsidated in an AAV particle. In particular, it can be encapsidated in an AAV1 particle, an AAV2 particle, an AAV3 particle, an AAV4 particle, an AAV5 particle or an AAV6 particle, a portion of any of these capsids, or a chimeric capsid particle as  
30 described above, by standard methods using the appropriate capsid proteins in the encapsidation process, as long as the nucleic acid vector fits within the size limitation

of the particle utilized. The encapsidation process itself is standard in the art. The AAV5 replication machinery, i.e. the rep initiator proteins and other functions required for replication, can be utilized to produce the AAV5 genome that can be packaged in an AAV1, 2, 3, 4, 5 or 6 capsid.

5

The recombinant AAV5 virion containing a vector can also be produced by recombinant methods utilizing multiple plasmids. In one example, the AAV5 rep nucleic acid would be cloned into one plasmid, the AAV5 ITR nucleic acid would be cloned into another plasmid and the AAV1, 2, 3, 4, 5 or 6 capsid nucleic acid would be  
10 cloned on another plasmid. These plasmids would then be introduced into cells. The cells that were efficiently transduced by all three plasmids, would exhibit specific integration as well as the ability to produce recombinant AAV5 virion. Additionally, two plasmids could be used where the AAV5 rep nucleic acid would be cloned into one plasmid and the AAV5 ITR and AAV5 capsid would be cloned into another plasmid.  
15 These plasmids would then be introduced into cells. The cells that were efficiently transduced by both plasmids, would exhibit specific integration as well as the ability to produce recombinant AAV5 virion.

An AAV5 capsid polypeptide encoding the entire VP1, VP2, and VP3  
20 polypeptide can have greater than 56% overall homology to the polypeptide having the amino acid sequence encoded by nucleotides in SEQ ID NOS:7,8 and 9, as shown in figures 4 and 5. The capsid protein can have about 70% homology, about 75% homology, 80% homology, 85% homology, 90% homology, 95% homology, 98% homology, 99% homology, or even 100% homology to the protein having the amino  
25 acid sequence encoded by the nucleotides set forth in SEQ ID NOS:7, 8 or 9. The percent homology used to identify proteins herein, can be based on a nucleotide-by-nucleotide comparison or more preferable is based on a computerized algorithm as described herein. Variations in the amino acid sequence of the AAV5 capsid protein are contemplated herein, as long as the resulting particle comprising an AAV5 capsid  
30 protein remains antigenically or immunologically distinct from AAV1, AAV2, AAV3, AAV4 or AAV6 capsid, as can be routinely determined by standard methods.

Specifically, for example, ELISA and Western blots can be used to determine whether a viral particle is antigenically or immunologically distinct from AAV2 or the other serotypes. Furthermore, the AAV5 particle preferably retains tissue tropism distinction from AAV2, such as that exemplified in the examples herein. An AAV5 chimeric  
5 particle comprising at least one AAV5 coat protein may have a different tissue tropism from that of an AAV5 particle consisting only of AAV5 coat proteins, but is still distinct from the tropism of an AAV2 particle, in that it will infect some cells not infected by AAV2 or an AAV2 particle.

10 The invention further provides a recombinant AAV5 virion, comprising an AAV5 particle containing, *i.e.*, encapsidating, a vector comprising a pair of AAV5 inverted terminal repeats. The recombinant vector can further comprise an AAV5 Rep-encoding nucleic acid. The vector encapsidated in the particle can further comprise an exogenous nucleic acid inserted between the inverted terminal repeats. AAV5 Rep  
15 confers targeted integration and efficient replication, thus production of recombinant AAV5, comprising AAV5 Rep, yields more particles than production of recombinant AAV2. Since AAV5 is more efficient at replicating and packaging its genome, the exogenous nucleic acid inserted, or in the AAV5 capsids of the present invention, between the inverted terminal repeats can be packaged in the AAV1, 2, 3, 4, or 6  
20 capsids to achieve the specific tissue tropism conferred by the capsid proteins.

The invention further contemplates chimeric recombinant ITRs that contains a rep binding site and a TRS site recognized by that Rep protein. By "Rep protein" is  
25 meant all four of the Rep proteins, Rep 40, Rep 78, Rep 52, Rep 68. Alternatively, "Rep protein" could be one or more of the Rep proteins described herein. One example of a chimeric ITR would consist of an AAV5 D region (SEQ ID NO: 23), an AAV5 TRS site (SEQ ID NO: 21), an AAV2 hairpin and an AAV2 binding site. Another example would be an AAV5 D region, an AAV5 TRS site, an AAV3 hairpin and an  
30 AAV3 binding site. In these chimeric ITRs, the D region can be from AAV1, 2, 3, 4, 5 or 6. The hairpin can be derived from AAV 1,2 3, 4, 5, 6. The binding site can be

derived from any of AAV1, 2, 3, 4, 5 or 6. Preferably, the D region and the TRS are from the same serotype.

The chimeric ITRs can be combined with AAV5 Rep protein and any of the  
5 AAV serotype capsids to obtain recombinant virion. For example, recombinant virion can be produced by an AAV5 D region, an AAV5 TRS site, an AAV2 hairpin, an AAV2 binding site, AAV5 Rep protein and AAV1 capsid. This recombinant virion would possess the cellular tropism conferred by the AAV1 capsid protein and would possess the efficient replication conferred by the AAV5 Rep.

10

Other examples of the ITR, Rep protein and Capsids that will produce recombinant virion are provided in the list below:

5ITR + 5Rep + 5Cap=virion  
15 5ITR + 5Rep + 1Cap=virion  
5ITR + 5Rep + 2Cap=virion  
5ITR + 5Rep + 3Cap=virion  
5ITR + 5Rep + 4Cap=virion  
5ITR + 5Rep + 6Cap=virion  
20 1ITR + 1Rep + 5Cap=virion  
2ITR + 2Rep + 5Cap=virion  
3ITR + 3Rep + 5Cap=virion  
4ITR + 4Rep + 5Cap=virion  
6ITR + 6Rep + 5Cap=virion

25

In any of the constructs described herein, inclusion of a promoter is preferred. As used in the constructs herein, unless otherwise specified, Cap (capsid) refers to any of AAV5 VP1, AAV5 VP2, AAV5 VP3, combinations thereof, functional fragments of any of VP1, VP2 or VP3, or chimeric capsids as described herein. The ITRs of the  
30 constructs described herein, can be chimeric recombinant ITRs as described elsewhere in the application.

Conjugates of recombinant or wild-type AAV5 virions and nucleic acids or proteins can be used to deliver those molecules to a cell. For example, the purified AAV5 can be used as a vehicle for delivering DNA bound to the exterior of the virus. Examples of this are to conjugate the DNA to the virion by a bridge using poly-L-lysine or other charged molecule. Also contemplated are virosomes that contain AAV5 structural proteins (AAV5 capsid proteins), lipids such as DOTAP, and nucleic acids that are complexed via charge interaction to introduce DNA into cells.

Also provided by this invention are conjugates that utilize the AAV5 capsid or a unique region of the AAV5 capsid protein (e.g. VP1, VP2 or VP3 or combinations thereof) to introduce DNA into cells. For example, the type 5 VP3 protein or fragment thereof, can be conjugated to a DNA on a plasmid that is conjugated to a lipid. Cells can be infected using the targeting ability of the VP3 capsid protein to achieve the desired tissue tropism, specific to AAV5. Type 5 VP1 and VP2 proteins can also be utilized to introduce DNA or other molecules into cells. By further incorporating the Rep protein and the AAV TRS into the DNA-containing conjugate, cells can be transduced and targeted integration can be achieved. For example, if AAV5 specific targeted integration is desired, a conjugate composed of the AAV5 VP3 capsid, AAV5 rep or a fragment of AAV5 rep, AAV5 TRS, the rep binding site, the heterologous DNA of interest, and a lipid, can be utilized to achieve AAV5 specific tropism and AAV5 specific targeted integration in the genome.

Further provided by this invention are chimeric viruses where AAV5 can be combined with herpes virus, herpes virus amplicons, baculovirus or other viruses to achieve a desired tropism associated with another virus. For example, the AAV5 ITRs could be inserted in the herpes virus and cells could be infected. Post-infection, the ITRs of AAV5 could be acted on by AAV5 rep provided in the system or in a separate vehicle to rescue AAV5 from the genome. Therefore, the cellular tropism of the herpes simplex virus can be combined with AAV5 rep mediated targeted integration. Other viruses that could be utilized to construct chimeric viruses include, lentivirus,

retrovirus, pseudotyped retroviral vectors, and adenoviral vectors.

The present invention further provides isolated nucleic acids of AAV5. For example, provided is an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:1 (AAV5 genome). This nucleic acid, or portions thereof, can be inserted into vectors, such as plasmids, yeast artificial chromosomes, or other viral vector (particle), if desired, by standard cloning methods. The present invention also provides an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:1. The nucleotides of SEQ ID NO:1 can have minor modifications and still be contemplated by the present invention. For example, modifications that do not alter the amino acid encoded by any given codon (such as by modification of the third, "wobble," position in a codon) can readily be made, and such alterations are known in the art. Furthermore, modifications that cause a resulting neutral (conserved) amino acid substitution of a similar amino acid can be made in a coding region of the genome. Additionally, modifications as described herein for the AAV5 components, such as the ITRs, the p5 promoter, etc. are contemplated in this invention. Furthermore, modifications to regions of SEQ ID NO:1 other than in the ITR, TRS Rep binding site and hairpin are likely to be tolerated without serious impact on the function of the nucleic acid as a recombinant vector.

20

As used herein, the term "isolated" refers to a nucleic acid separated or significantly free from at least some of the other components of the naturally occurring organism, for example, the cell structural components or viral components commonly found associated with nucleic acids in the environment of the virus and/or other nucleic acids. The isolation of the native nucleic acids can be accomplished, for example, by techniques such as cell lysis followed by phenol plus chloroform extraction, followed by ethanol precipitation of the nucleic acids. The nucleic acids of this invention can be isolated from cells according to any of many methods well known in the art.

30

As used herein, the term "nucleic acid" refers to single-or multiple stranded molecules which may be DNA or RNA, or any combination thereof, including

modifications to those nucleic acids. The nucleic acid may represent a coding strand or its complement, or any combination thereof. Nucleic acids may be identical in sequence to the sequences which are naturally occurring for any of the novel genes discussed herein or may include alternative codons which encode the same amino acid as those provided herein, including that which is found in the naturally occurring sequence. These nucleic acids can also be modified from their typical structure. Such modifications include, but are not limited to, methylated nucleic acids, the substitution of a non-bridging oxygen on the phosphate residue with either a sulfur (yielding phosphorothioate deoxynucleotides), selenium (yielding phosphorselenoate deoxynucleotides), or methyl groups (yielding methylphosphonate deoxynucleotides).

The present invention additionally provides an isolated nucleic acid that selectively hybridizes with any nucleic acid disclosed herein, including the entire AAV5 genome and any unique fragment thereof, including the Rep and capsid encoding sequences (e.g. SEQ ID NOS: 1, 7, 8, 9, 10, 11, 13, 15, 16, 17, 18, 19, 20, 21, 22 and 23). Specifically, the nucleic acid can selectively or specifically hybridize to an isolated nucleic acid consisting of the nucleotide sequence set forth in SEQ ID NO:1 (AAV5 genome). The present invention further provides an isolated nucleic acid that selectively or specifically hybridizes with an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:1 (AAV5 genome). By "selectively hybridizes" as used herein is meant a nucleic acid that hybridizes to one of the disclosed nucleic acids under sufficient stringency conditions without significant hybridization to a nucleic acid encoding an unrelated protein, and particularly, without detectably hybridizing to nucleic acids of AAV2. Thus, a nucleic acid that selectively hybridizes with a nucleic acid of the present invention will not selectively hybridize under stringent conditions with a nucleic acid encoding a different protein or the corresponding protein from a different serotype of the virus, and vice versa. A "specifically hybridizing" nucleic acid is one that hybridizes under stringent conditions to only a nucleic acid found in AAV5. Therefore, nucleic acids for use, for example, as primers and probes to detect or amplify the target nucleic acids are contemplated herein. Nucleic acid fragments that selectively hybridize to any given nucleic acid can

be used, *e.g.*, as primers and or probes for further hybridization or for amplification methods (*e.g.*, polymerase chain reaction (PCR), ligase chain reaction (LCR)).

Additionally, for example, a primer or probe can be designed that selectively hybridizes with both AAV5 and a gene of interest carried within the AAV5 vector (*i.e.*, a chimeric  
5 nucleic acid).

Stringency of hybridization is controlled by both temperature and salt concentration of either or both of the hybridization and washing steps. Typically, the stringency of hybridization to achieve selective hybridization involves hybridization in  
10 high ionic strength solution (6X SSC or 6X SSPE) at a temperature that is about 12-25°C below the  $T_m$  (the melting temperature at which half of the molecules dissociate from their hybridization partners) followed by washing at a combination of temperature and salt concentration chosen so that the washing temperature is about 5°C to 20°C below the  $T_m$ . The temperature and salt conditions are readily determined empirically  
15 in preliminary experiments in which samples of reference DNA immobilized on filters are hybridized to a labeled nucleic acid of interest and then washed under conditions of different stringencies. Hybridization temperatures are typically higher for DNA-RNA and RNA-RNA hybridizations. The washing temperatures can be used as described above to achieve selective stringency, as is known in the art. (Sambrook et al.,  
20 *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989; Kunkel et al. *Methods Enzymol.* 1987:154:367, 1987). A preferable stringent hybridization condition for a DNA:DNA hybridization can be at about 68°C (in aqueous solution) in 6X SSC or 6X SSPE followed by washing at 68°C. Stringency of hybridization and washing, if desired, can be reduced  
25 accordingly as the degree of complementarity desired is decreased, and further, depending upon the G-C or A-T richness of any area wherein variability is searched for. Likewise, stringency of hybridization and washing, if desired, can be increased accordingly as homology desired is increased, and further, depending upon the G-C or A-T richness of any area wherein high homology is desired, all as known in the art.



A nucleic acid that selectively hybridizes to any portion of the AAV5 genome is contemplated herein. Therefore, a nucleic acid that selectively hybridizes to AAV5 can be of longer length than the AAV5 genome, it can be about the same length as the AAV5 genome or it can be shorter than the AAV5 genome. The length of the nucleic acid is limited on the shorter end of the size range only by its specificity for hybridization to AAV5, *i.e.*, once it is too short, typically less than about 5 to 7 nucleotides in length, it will no longer bind specifically to AAV5, but rather will hybridize to numerous background nucleic acids. Additionally contemplated by this invention is a nucleic acid that has a portion that specifically hybridizes to AAV5 and a portion that specifically hybridizes to a gene of interest inserted within AAV5.

The present invention further provides an isolated nucleic acid encoding an adeno-associated virus 5 Rep protein. The AAV5 Rep proteins are encoded by open reading frame (ORF) 1 of the AAV5 genome. Examples of the AAV5 Rep genes are shown in the nucleic acid set forth in SEQ ID NO:1, and include nucleic acids consisting essentially of the nucleotide sequences set forth in SEQ ID NOS:10 (Rep52), 11 (Rep78), 13 (Rep40), and 15 (Rep68), and nucleic acids comprising the nucleotide sequences set forth in SEQ ID NOS:10, 11, 13, and 15. However, the present invention contemplates that the Rep nucleic acid can include any one, two, three, or four of the four Rep proteins, in any order, in such a nucleic acid. Furthermore, minor modifications are contemplated in the nucleic acid, such as silent mutations in the coding sequences, mutations that make neutral or conservative changes in the encoded amino acid sequence, and mutations in regulatory regions that do not disrupt the expression of the gene. Examples of other minor modifications are known in the art. Further modifications can be made in the nucleic acid, such as to disrupt or alter expression of one or more of the Rep proteins in order to, for example, determine the effect of such a disruption; such as to mutate one or more of the Rep proteins to determine the resulting effect, etc. However, in general, a modified nucleic acid encoding a Rep protein will have at least about 85%, about 90%, about 93%, about 95%, about 98% or 100% homology to the Rep nucleic sequences described herein *e.g.*, SEQ ID NOS: 10, 11, 13 and 15, and the Rep polypeptide encoded therein will

have overall about 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence described herein, e.g., SEQ ID NOS:2, 3, 12 and 14. Percent homology is determined by the techniques described herein.

5           The present invention also provides an isolated nucleic acid that selectively or specifically hybridizes with a nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NOS:10, 11, 13 and 15, and an isolated nucleic acid that selectively hybridizes with a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NOS:10, 11, 13 and 15. "Selectively hybridizing" and "stringency of  
10 hybridization" is defined elsewhere herein.

As described above, the present invention provides the nucleic acid encoding a Rep 40 protein and, in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 13, an isolated nucleic acid consisting essentially of  
15 the nucleotide sequence set forth in SEQ ID NO: 13, and a nucleic acid encoding the adeno-associated virus 5 protein having the amino acid sequence set forth in SEQ ID NO: 12. The present invention also provides the nucleic acid encoding a Rep 52 protein, and in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:10, an isolated nucleic acid consisting essentially of the  
20 nucleotide sequence set forth in SEQ ID NO:10, and a nucleic acid encoding the adeno-associated virus 5 Rep protein having the amino acid sequence set forth in SEQ ID NO:2. The present invention further provides the nucleic acid encoding a Rep 68 protein and, in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 15, an isolated nucleic acid consisting essentially of the  
25 nucleotide sequence set forth in SEQ ID NO: 15, and a nucleic acid encoding the adeno-associated virus 5 protein having the amino acid sequence set forth in SEQ ID NO: 14. And, further, the present invention provides the nucleic acid encoding a Rep 78 protein, and in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:11, an isolated nucleic acid consisting essentially of  
30 the nucleotide sequence set forth in SEQ ID NO:11, and a nucleic acid encoding the adeno-associated virus 5 Rep protein having the amino acid sequence set forth in SEQ

ID NO:3. As described elsewhere herein, these nucleic acids can have minor modifications, including silent nucleotide substitutions, mutations causing conservative amino acid substitutions in the encoded proteins, and mutations in control regions that do not or minimally affect the encoded amino acid sequence.

5

The present invention further provides a nucleic acid encoding the entire AAV5 Capsid polypeptide. Furthermore, the present invention provides a nucleic acid encoding each of the three AAV5 coat proteins, VP1, VP2, and VP3. Thus, the present invention provides a nucleic acid encoding AAV5 VP1, a nucleic acid encoding AAV5  
10 VP2, and a nucleic acid encoding AAV5 VP3. Thus, the present invention provides a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO:4 (VP1); a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO:5 (VP2), and a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO:6 (VP3). The present invention also specifically provides a nucleic acid comprising SEQ ID NO:7  
15 (VP1 gene); a nucleic acid comprising SEQ ID NO:8 (VP2 gene); and a nucleic acid comprising SEQ ID NO:9 (VP3 gene). The present invention also specifically provides a nucleic acid consisting essentially of SEQ ID NO:7 (VP1 gene), a nucleic acid consisting essentially of SEQ ID NO:8 (VP2 gene), and a nucleic acid consisting essentially of SEQ ID NO:9 (VP3 gene). Minor modifications in the nucleotide  
20 sequences encoding the capsid, or coat, proteins are contemplated, as described above for other AAV5 nucleic acids. However, in general, a modified nucleic acid encoding a capsid protein will have at least about 85%, about 90%, about 93%, about 95%, about 98% or 100% homology to the capsid nucleic sequences described herein e.g., SEQ ID NOS: 7, 8, and 9, and the capsid polypeptide encoded therein will have overall about  
25 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence described herein, e.g., SEQ ID NOS:4, 5, and 6. Nucleic acids that selectively hybridize with the nucleic acids of SEQ ID NOS:7,8 and 9 under the conditions described above are also provided.

30

The present invention also provides a cell containing one or more of the herein described nucleic acids, such as the AAV5 genome, AAV5 ORF1 and ORF2, each

AAV5 Rep protein gene, or each AAV5 capsid protein gene. Such a cell can be any desired cell and can be selected based upon the use intended. For example, cells can include bacterial cells, yeast cells, insect cells, human HeLa cells and simian Cos cells as well as other human and mammalian cells and cell lines. Primary cultures as well as  
5 established cultures and cell lines can be used. Nucleic acids of the present invention can be delivered into cells by any selected means, in particular depending upon the target cells. Many delivery means are well-known in the art. For example, electroporation, calcium phosphate precipitation, microinjection, cationic or anionic liposomes, and liposomes in combination with a nuclear localization signal peptide for  
10 delivery to the nucleus can be utilized, as is known in the art. Additionally, if the nucleic acids are in a viral particle, the cells can simply be transduced with the virion by standard means known in the art for AAV transduction. Small amounts of the recombinant AAV5 virus can be made to infect cells and produce more of itself.

15       The invention provides purified AAV5 polypeptides. The term "polypeptide" as used herein refers to a polymer of amino acids and includes full-length proteins and fragments thereof. Thus, "protein," polypeptide," and "peptide" are often used interchangeably herein. Substitutions can be selected by known parameters to be neutral (*see, e.g.,* Robinson WE Jr, and Mitchell WM., AIDS 4:S151-S162 (1990)).  
20 As will be appreciated by those skilled in the art, the invention also includes those polypeptides having slight variations in amino acid sequences or other properties. Such variations may arise naturally as allelic variations (*e.g.,* due to genetic polymorphism) or may be produced by human intervention (*e.g.,* by mutagenesis of cloned DNA sequences), such as induced point, deletion, insertion and substitution mutants. Minor  
25 changes in amino acid sequence are generally preferred, such as conservative amino acid replacements, small internal deletions or insertions, and additions or deletions at the ends of the molecules. Substitutions may be designed based on, for example, the model of Dayhoff, *et al.* (in *Atlas of Protein Sequence and Structure* 1978, Nat'l Biomed. Res. Found., Washington, D.C.). These modifications can result in changes in  
30 the amino acid sequence, provide silent mutations, modify a restriction site, or provide other specific mutations. The location of any modifications to the polypeptide will

often determine its impact on function. Particularly, alterations in regions non-essential to protein function will be tolerated with fewer effects on function. Elsewhere in the application regions of the AAV5 proteins are described to provide guidance as to where substitutions, additions or deletions can be made to minimize the likelihood of  
5 disturbing the function of the variant.

A polypeptide of the present invention can be readily obtained by any of several means. For example, the polypeptide of interest can be synthesized chemically by standard methods. Additionally, the coding regions of the genes can be recombinantly  
10 expressed and the resulting polypeptide isolated by standard methods. Furthermore, an antibody specific for the resulting polypeptide can be raised by standard methods (see, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1988), and the protein can be isolated from a cell expressing the nucleic acid encoding the polypeptide by selective  
15 hybridization with the antibody. This protein can be purified to the extent desired by standard methods of protein purification (see, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989).

20 Typically, to be unique, a polypeptide fragment of the present invention will be at least about 5 amino acids in length; however, unique fragments can be 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 or more amino acids in length. A unique polypeptide will typically comprise such a unique fragment; however, a unique polypeptide can also be determined by its overall homology. A unique polypeptide can be 6, 7, 8, 9, 10, 20,  
25 30, 40, 50, 60, 70, 80, 90, 100 or more amino acids in length. Uniqueness of a polypeptide fragment can readily be determined by standard methods such as searches of computer databases of known peptide or nucleic acid sequences or by hybridization studies to the nucleic acid encoding the protein or to the protein itself, as known in the art. The uniqueness of a polypeptide fragment can also be determined immunologically  
30 as well as functionally. Uniqueness can be simply determined in an amino acid-by-amino acid comparison of the polypeptides.

An antigenic or immunoreactive fragment of this invention is typically an amino acid sequence of at least about 5 consecutive amino acids, and it can be derived from the AAV5 polypeptide amino acid sequence. An antigenic AAV5 fragment is any fragment unique to the AAV5 protein, as described herein, against which an AAV5-specific antibody can be raised, by standard methods. Thus, the resulting antibody-antigen reaction should be specific for AAV5.

The present invention provides an isolated AAV5 Rep protein. An AAV5 Rep polypeptide is encoded by ORF1 of AAV5. The present invention also provides each individual AAV5 Rep protein. Thus the present invention provides AAV5 Rep 40 (e.g., SEQ ID NO: 12), or a unique fragment thereof. The present invention provides AAV5 Rep 52 (e.g., SEQ ID NO: 2), or a unique fragment thereof. The present invention provides AAV5 Rep 68 (e.g., SEQ ID NO: 14), or a unique fragment thereof. The present invention provides an example of AAV5 Rep 78 (e.g., SEQ ID NO: 3), or a unique fragment thereof. By "unique fragment thereof" is meant any smaller polypeptide fragment encoded by an AAV5 rep gene that is of sufficient length to be found only in the Rep polypeptide. Substitutions and modifications of the amino acid sequence can be made as described above and, further, can include protein processing modifications, such as glycosylation, to the polypeptide.

20

The present invention further provides an AAV5 Capsid polypeptide or a unique fragment thereof. AAV5 capsid polypeptide is encoded by ORF 2 of AAV5. The present invention further provides the individual AAV5 capsid proteins, VP1, VP2 and VP3 or unique fragments thereof. Thus, the present invention provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:4 (VP1). The present invention additionally provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:5 (VP2). The present invention also provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:6 (VP3). By "unique fragment thereof" is meant any smaller polypeptide fragment encoded by any AAV5 capsid gene that is of sufficient length to be found only in the AAV5 capsid protein. Substitutions and modifications of the amino acid sequence can be made as

25  
30

described above and, further, can include protein processing modifications, such as glycosylation, to the polypeptide. However, an AAV5 Capsid polypeptide including all three coat proteins will have greater than about 56% overall homology to the polypeptide encoded by the nucleotides set forth in SEQ ID NOS:4,5 or 6. The protein  
5 can have about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, 93%, 95%, 97% or even 100% homology to the amino acid sequence encoded by the nucleotides set forth in SEQ ID NOS:4,5 or 6. An AAV5 VP1 polypeptide can have at least about 58%, about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or about 100% homology to the amino acid sequence set forth in SEQ ID NO:4. An  
10 AAV5 VP2 polypeptide can have at least about 58%, about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or about 100% homology to the amino acid sequence set forth in SEQ ID NO:5. An AAV5 VP3 polypeptide can have at least about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or about 100% homology to the amino acid sequence set forth in SEQ ID NO:6.

15

The present invention further provides an isolated antibody that specifically binds an AAV5 Rep protein or a unique epitope thereof. Also provided are isolated antibodies that specifically bind the AAV5 Rep 52 protein, the AAV5 Rep 40 protein, the AAV5 Rep 68 protein and the AAV5 Rep 78 protein having the amino acid  
20 sequences set forth in SEQ ID NO:2, SEQ ID NO: 12, SEQ ID NO: 14 and SEQ ID NO: 3, respectively or that specifically binds a unique fragment thereof. Clearly, any given antibody can recognize and bind one of a number of possible epitopes present in the polypeptide; thus only a unique portion of a polypeptide (having the epitope) may need to be present in an assay to determine if the antibody specifically binds the  
25 polypeptide.

The present invention additionally provides an isolated antibody that specifically binds any of the adeno-associated virus 5 Capsid proteins (VP1, VP2 or VP3), a unique epitope thereof, or the polypeptide comprising all three AAV5 coat  
30 proteins. Also provided is an isolated antibody that specifically binds the AAV5 capsid protein having the amino acid sequence set forth in SEQ ID NO:4 (VP1), or that

specifically binds a unique fragment thereof. The present invention further provides an isolated antibody that specifically binds the AAV5 Capsid protein having the amino acid sequence set forth in SEQ ID NO:5 (VP2), or that specifically binds a unique fragment thereof. The invention additionally provides an isolated antibody that

5 specifically binds the AAV5 Capsid protein having the amino acid sequence set forth in SEQ ID NO:6 (VP3), or that specifically binds a unique fragment thereof. Again, any given antibody can recognize and bind one of a number of possible epitopes present in the polypeptide; thus only a unique portion of a polypeptide (having the epitope) may need to be present in an assay to determine if the antibody specifically binds the

10 polypeptide.

The antibody can be a component of a composition that comprises an antibody that specifically binds the AAV5 protein. The composition can further comprise, *e.g.*, serum, serum-free medium, or a pharmaceutically acceptable carrier such as

15 physiological saline, etc..

By "an antibody that specifically binds" an AAV5 polypeptide or protein is meant an antibody that selectively binds to an epitope on any portion of the AAV5 peptide such that the antibody binds specifically to the corresponding AAV5

20 polypeptide without significant background. Specific binding by an antibody further means that the antibody can be used to selectively remove the target polypeptide from a sample comprising the polypeptide or and can readily be determined by radioimmunoassay (RIA), bioassay, or enzyme-linked immunosorbant (ELISA) technology. An ELISA method effective for the detection of the specific antibody-antigen binding can, for example, be as follows: (1) bind the antibody to a substrate;

25 (2) contact the bound antibody with a sample containing the antigen; (3) contact the above with a secondary antibody bound to a detectable moiety (*e.g.*, horseradish peroxidase enzyme or alkaline phosphatase enzyme); (4) contact the above with the substrate for the enzyme; (5) contact the above with a color reagent; (6) observe the

30 color change.



An antibody can include antibody fragments such as Fab fragments which retain the binding activity. Antibodies can be made as described in, *e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York (1988). Briefly, purified antigen can be injected into an animal in  
5 an amount and in intervals sufficient to elicit an immune response. Antibodies can either be purified directly, or spleen cells can be obtained from the animal. The cells are then fused with an immortal cell line and screened for antibody secretion. Individual hybridomas are then propagated as individual clones serving as a source for a particular monoclonal antibody.

10

The present invention additionally provides a method of screening a cell for infectivity by AAV5 comprising contacting the cell with AAV5 and detecting the presence of AAV5 in the cells. AAV5 particles can be detected using any standard physical or biochemical methods. For example, physical methods that can be used for  
15 this detection include DNA based methods such as 1) polymerase chain reaction (PCR) for viral DNA or RNA or 2) direct hybridization with labeled probes, and immunological methods such as by 3) antibody directed against the viral structural or non- structural proteins. Catalytic methods of viral detection include, but are not limited to, detection of site and strand specific DNA nicking activity of Rep proteins or  
20 replication of an AAV origin- containing substrate. Reporter genes can also be utilized to detect cells that transduce AAV-5. For example,  $\beta$ -gal, green fluorescent protein or luciferase can be inserted into a recombinant AAV-5. The cell can then be contacted with the recombinant AAV-5, either *in vitro* or *in vivo* and a colorimetric assay could detect a color change in the cells that would indicate transduction of AAV-5 in the cell.  
25 Additional detection methods are outlined in Fields, *Virology*, Raven Press, New York, New York. 1996.

For screening a cell for infectivity by AAV5, wherein the presence of AAV5 in the cells is determined by nucleic acid hybridization methods, a nucleic acid probe for  
30 such detection can comprise, for example, a unique fragment of any of the AAV5 nucleic acids provided herein. The uniqueness of any nucleic acid probe can readily be

determined as described herein. Additionally, the presence of AAV5 in cells can be determined by fluorescence, antibodies to gene products, focus forming assays, plaque lifts, Western blots and chromogenic assays. The nucleic acid can be, for example, the nucleic acid whose nucleotide sequence is set forth in SEQ ID NO: 1, 7, 8, 9, 10, 11, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23 or a unique fragment thereof.

The present invention includes a method of determining the suitability of an AAV5 vector for administration to a subject comprising administering to an antibody-containing sample from the subject an antigenic fragment of an isolated AAV5 Rep or Capsid protein, and detecting neutralizing antibody-antigen reaction in the sample, the presence of a neutralizing reaction indicating the AAV5 vector may be unsuitable for use in the subject. The present method of determining the suitability of an AAV5 vector for administration to a subject can comprise contacting an antibody-containing sample from the subject with a unique antigenic or immunogenic fragment of an AAV5 Rep protein (e.g. Rep 40, Rep 52, Rep 68, Rep 78) and detecting an antibody-antigen reaction in the sample, the presence of a reaction indicating the AAV5 vector to be unsuitable for use in the subject. The AAV5 Rep proteins are provided herein, and their antigenic fragments are routinely determined. The AAV5 capsid protein can be used to select an antigenic or immunogenic fragment, for example from the amino acid sequence set forth in SEQ ID NO:4 (VP1), the amino acid sequence set forth in SEQ ID NO: 5 (VP2) or the amino acid sequence set forth in SEQ ID NO:6 (VP3). Alternatively, or additionally, an antigenic or immunogenic fragment of an isolated AAV5 Rep protein can be utilized in this determination method. The AAV5 Rep protein from which an antigenic fragment is selected can have the amino acid sequence encoded by the nucleic acid set forth in SEQ ID NO:1, the amino acid sequence set forth in SEQ ID NO:2, or the amino acid sequence set forth in SEQ ID NO:3, the amino acid sequence set forth in SEQ ID NO: 12, or the amino acid sequence set forth in SEQ ID NO:14.

The AAV5 polypeptide fragments can be analyzed to determine their antigenicity, immunogenicity and/or specificity. Briefly, various concentrations of a putative immunogenically specific fragment are prepared and administered to a subject and the immunological response (e.g., the production of antibodies or cell mediated immunity) of an animal to each concentration is determined. The amounts of antigen administered depend on the subject, e.g. a human, rabbit or a guinea pig, the condition of the subject, the size of the subject, etc. Thereafter an animal so inoculated with the antigen can be exposed to the AAV5 viral particle or AAV5 protein to test the immunoreactivity or the antigenicity of the specific immunogenic fragment. The specificity of a putative antigenic or immunogenic fragment can be ascertained by testing sera, other fluids or lymphocytes from the inoculated animal for cross reactivity with other closely related viruses, such as AAV1, AAV2, AAV3, AAV4 and AAV5.

The hemagglutination assay can also be used to rapidly identify and detect AAV5 viral particles. Detection of hemagglutination activity correlates with infectivity and can be used to titer the virus. This assay could also be used to identify antibodies in a patients serum which might interact with the virus. Hemagglutination has been shown to correlate with infectivity and therefore hemagglutination may be a useful assay for identify cellular receptors for AAV5.

20

By the "suitability of an AAV5 vector for administration to a subject" is meant a determination of whether the AAV5 vector will elicit a neutralizing immune response upon administration to a particular subject. A vector that does not elicit a significant immune response is a potentially suitable vector, whereas a vector that elicits a significant, neutralizing immune response (e.g. at least 90%) is thus likely to be unsuitable for use in that subject. Significance of any detectable immune response is a standard parameter understood by the skilled artisan in the field. For example, one can incubate the subject's serum with the virus, then determine whether that virus retains its ability to transduce cells in culture. If such virus cannot transduce cells in culture, the vector likely has elicited a significant immune response.

30

Alternatively, or additionally, one skilled in the art could determine whether or not AAV5 administration would be suitable for a particular cell type of a subject. For example, the artisan could culture muscle cells *in vitro* and transduce the cells with AAV5 in the presence or absence of the subject's serum. If there is a reduction in  
5 transduction efficiency, this could indicate the presence of a neutralizing antibody or other factors that may inhibit transduction. Normally, greater than 90% inhibition would have to be observed in order to rule out the use of AAV-5 as a vector. However, this limitation could be overcome by treating the subject with an immunosuppressant that could block the factors inhibiting transduction.

10

As will be recognized by those skilled in the art, numerous types of immunoassays are available for use in the present invention to detect binding between an antibody and an AAV5 polypeptide of this invention. For instance, direct and indirect binding assays, competitive assays, sandwich assays, and the like, as are  
15 generally described in, *e.g.*, U.S. Pat. Nos. 4,642,285; 4,376,110; 4,016,043; 3,879,262; 3,852,157; 3,850,752; 3,839,153; 3,791,932; and Harlow and Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, N.Y. (1988). For example, enzyme immunoassays such as immunofluorescence assays (IFA), enzyme linked immunosorbent assays (ELISA) and immunoblotting can be readily adapted to  
20 accomplish the detection of the antibody. An ELISA method effective for the detection of the antibody bound to the antigen can, for example, be as follows: (1) bind the antigen to a substrate; (2) contact the bound antigen with a fluid or tissue sample containing the antibody; (3) contact the above with a secondary antibody specific for the antigen and bound to a detectable moiety (*e.g.*, horseradish peroxidase enzyme or  
25 alkaline phosphatase enzyme); (4) contact the above with the substrate for the enzyme; (5) contact the above with a color reagent; (6) observe color change.

The antibody-containing sample of this method can comprise any biological sample which would contain the antibody or a cell containing the antibody, such as  
30 blood, plasma, serum, bone marrow, saliva and urine.

The present invention also provides a method of producing the AAV5 virus by transducing a cell with the nucleic acid encoding the virus.

The present method further provides a method of delivering an exogenous  
5 (heterologous) nucleic acid to a cell comprising administering to the cell an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to the cell.

10 The AAV ITRs in the vector for the herein described delivery methods can be AAV5 ITRs (SEQ ID NOS: 19 and 20). Furthermore, the AAV ITRs in the vector for the herein described nucleic acid delivery methods can also comprise AAV1, AAV2, AAV3, AAV4, or AAV6 inverted terminal repeats.

15 The present invention also includes a method of delivering a heterologous nucleic acid to a subject comprising administering to a cell from the subject an AAV5 virion or particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, and returning the cell to the subject, thereby delivering the nucleic acid to the subject. The AAV ITRs can be any AAV ITRs, including AAV5 ITRs and  
20 AAV2 ITRs. For example, in an *ex vivo* administration, cells are isolated from a subject by standard means according to the cell type and placed in appropriate culture medium, again according to cell type (*see, e.g.,* ATCC catalog). Viral particles are then contacted with the cells as described above, and the virus is allowed to transduce the cells. Cells can then be transplanted back into the subject's body, again by means  
25 standard for the cell type and tissue (*e. g.,* in general, U.S. Patent No. 5,399,346; for neural cells, Dunnett, S.B. and Björklund, A., eds., *Transplantation: Neural Transplantation-A Practical Approach*, Oxford University Press, Oxford (1992)). If desired, prior to transplantation, the cells can be studied for degree of transduction by the virus, by known detection means and as described herein. Cells for *ex vivo*  
30 transduction followed by transplantation into a subject can be selected from those listed above, or can be any other selected cell including progenitor cells of the cells listed

above. Preferably, a selected cell type is examined for its capability to be transfected by AAV5. Preferably, the selected cell will be a cell readily transduced with AAV5 particles; however, depending upon the application, even cells with relatively low transduction efficiencies can be useful, particularly if the cell is from a tissue or organ in which even production of a small amount of the protein or antisense RNA encoded by the vector will be beneficial to the subject.

The present invention further provides a method of delivering a nucleic acid to a cell in a subject comprising administering to the subject an AAV5 virion or particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to a cell in the subject. Administration can be an *ex vivo* administration directly to a cell removed from a subject, such as any of the cells listed above, followed by replacement of the cell back into the subject, or administration can be *in vivo* administration to a cell in the subject. For *ex vivo* administration, cells are isolated from a subject by standard means according to the cell type and placed in appropriate culture medium, again according to cell type (*see, e.g.*, ATCC catalog). Viral particles are then contacted with the cells as described above, and the virus is allowed to transfect the cells. Cells can then be transplanted back into the subject's body, again by means standard for the cell type and tissue (*e. g.*, for neural cells, Dunnett, S.B. and Björklund, A., eds., *Transplantation: Neural Transplantation-A Practical Approach*, Oxford University Press, Oxford (1992)). If desired, prior to transplantation, the cells can be studied for degree of transfection by the virus, by known detection means and as described herein.

The present invention further provides a method of delivering a nucleic acid to a cell in a subject having neutralizing antibodies to AAV2 comprising administering to the subject an AAV5 virion or particle comprising the nucleic acid, thereby delivering the nucleic acid to a cell in the subject. A subject that has neutralizing antibodies to AAV2 can readily be determined by any of several known means, such as contacting AAV2 protein(s) with an antibody-containing sample, such as blood, from a subject and detecting an antigen-antibody reaction in the sample. Delivery of the AAV5

particle can be by either *ex vivo* or *in vivo* administration as herein described. Thus, a subject who might have an adverse immunogenic reaction to a vector administered in an AAV2 viral particle can have a desired nucleic acid delivered using an AAV5 particle. This delivery system can be particularly useful for subjects who have received therapy utilizing AAV2 particles in the past and have developed antibodies to AAV2. An AAV5 regimen can now be substituted to deliver the desired nucleic acid.

In any of the methods of delivering heterologous nucleic acids to a cell or subject described herein, the AAV5-conjugated nucleic acid or AAV5 particle-conjugated nucleic acids described herein can be used.

*In vivo* administration to a human subject or an animal model can be by any of many standard means for administering viruses, depending upon the target organ, tissue or cell. Virus particles can be administered orally, parenterally (*e.g.*, intravenously), by intramuscular injection, by direct tissue or organ injection, by intraperitoneal injection, topically, transdermally, via aerosol delivery, via the mucosa or the like. Viral nucleic acids (non-encapsidated) can also be administered, *e.g.*, as a complex with cationic liposomes, or encapsulated in anionic liposomes. The present compositions can include various amounts of the selected viral particle or non-encapsidated viral nucleic acid in combination with a pharmaceutically acceptable carrier and, in addition, if desired, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents, etc. Parental administration, if used, is generally characterized by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Dosages will depend upon the mode of administration, the disease or condition to be treated, and the individual subject's condition, but will be that dosage typical for and used in administration of other AAV vectors, such as AAV2 vectors. Often a single dose can be sufficient; however, the dose can be repeated if desirable.

Administration methods can be used to treat brain disorders such as Parkinson's disease, Alzheimer's disease, and demyelination disease. Other diseases that can be

treated by these methods include metabolic disorders such as , musculoskeletal diseases, cardiovascular disease, cancer, and autoimmune disorders.

Administration of this recombinant AAV5 virion or particle to the cell can be  
5 accomplished by any means, including simply contacting the particle, optionally  
contained in a desired liquid such as tissue culture medium, or a buffered saline  
solution, with the cells. The virion can be allowed to remain in contact with the cells  
for any desired length of time, and typically the virion is administered and allowed to  
remain indefinitely. For such *in vitro* methods, the virion can be administered to the  
10 cell by standard viral transduction methods, as known in the art and as exemplified  
herein. Titers of virus to administer can vary, particularly depending upon the cell type,  
but will be typical of that used for AAV transduction in general which is well known in  
the art. Additionally the titers used to transduce the particular cells in the present  
examples can be utilized.

15

The cells that can be transduced by the present recombinant AAV5 virion or  
particle can include any desired cell, such as the following cells and cells derived from  
the following tissues, human as well as other mammalian tissues, such as primate,  
horse, sheep, goat, pig, dog, rat, and mouse: Adipocytes, Adenocyte, Adrenal cortex,  
20 Airway epithelial cells, Alveolar cells, Amnion, Aorta, Ascites, Astrocyte, Bladder,  
Bone, Bone marrow, Brain, Breast, Bronchus, Cardiac muscle, Cecum, Cerebellar,  
Cervix, Chorion, Colon, Conjunctiva, Connective tissue, Cornea, Dermis, Duodenum,  
Endometrium, Endothelium, Endothelial cells, Ependymal cells, Epithelial tissue,  
Epithelial cells, Epidermis, Esophagus, Eye, Fascia, Fibroblasts, Foreskin, Gastric,  
25 Glial cells, Glioblast, Gonad, Hepatic cells, Histocyte, Ileum, Intestine, small Intestine,  
Jejunum, Keratinocytes, Kidney, Larynx, Leukocytes, Lipocyte, Liver, Lung, Lymph  
node, Lymphoblast, Lymphocytes, Macrophages, Mammary alveolar nodule, Mammary  
gland, Mastocyte, Maxilla, Melanocytes, Mesenchymal, Monocytes, Mouth, Myelin,  
Myoblasts Nervous tissue, Neuroblast, Neurons, Neuroglia, Osteoblasts, Osteogenic  
30 cells, Ovary, Palate, Pancreas, Papilloma, Peritoneum, Pituicytes, Pharynx, Placenta,  
Plasma cells, Pleura, Prostate, Rectum, Salivary gland, Skeletal muscle, Skin, Smooth



muscle, Somatic, Spinal cord, Spleen, Squamous, Stomach, Submandibular gland, Submaxillary gland, Synoviocytes, Testis, Thymus, Thyroid, Trabeculae, Trachea, Turbinate, Umbilical cord, Ureter, and Uterus. Thus, the particles and virions of the present invention can be used to deliver a nucleic acid to these cells.

5

More specifically, the present invention provides a method of delivering a nucleic acid to an ependymal cell, comprising administering to the ependymal cell an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to the ependymal

10 cell.

Also provided by the present invention is a method of delivering a nucleic acid to a neuron, comprising administering to the neuron an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal

15 repeats, thereby delivering the nucleic acid to the neuron.

Further provided by this invention is a method of delivering a nucleic acid to an astrocyte, comprising administering to an astrocyte an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal

20 repeats, thereby delivering the nucleic acid to an astrocyte.

The present invention also provides a method of delivering a nucleic acid to an airway epithelial cell, comprising administering to an airway epithelial cell an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV

25 inverted terminal repeats, thereby delivering the nucleic acid to the airway epithelial cell.

The present invention also provides a method of delivering a nucleic acid to an alveolar cell, comprising administering to an alveolar cell an AAV5 particle containing

30 a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to the alveolar cell.

The present invention also provides a method of delivering a nucleic acid to a cerebellar cell, comprising administering to a cerebellar cell an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to the cerebellar cell.

5

Also provided is a method of delivering a nucleic acid to an ependymal cell in a subject comprising administering to the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to an ependymal cell in the subject.

10

Further provided is a method of delivering a nucleic acid to a neuron in a subject comprising administering to the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to a neuron in the subject.

15

Also provided is a method of delivering a nucleic acid to an astrocyte in a subject comprising administering to the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to an astrocyte in the subject.

20

Also provided is a method of delivering a nucleic acid to an alveolar cell in a subject comprising administering to the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to an alveolar cell in the subject.

25

Also provided is a method of delivering a nucleic acid to a cerebellar cell in a subject comprising administering to the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to a cerebellar cell in the subject.

30

Further provided is a method of delivering a nucleic acid to an airway epithelial cell in a subject comprising administering to the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to an airway epithelial cell in the subject.

5

The use of AAV5 to deliver genes to the airway epithelia would be of benefit in genetic diseases like cystic fibrosis, pseudohypoaldosteronism, and immotile cilia syndrome. Furthermore, delivering genes to the airway epithelia would be of impact in several non-genetic diseases. For example, delivering genes that make antibiotic like peptides would be useful to prevent or treat bronchitis; delivering genes that make growth factors would be of value in common diseases like chronic bronchitis. Also, AAV5 could be used to deliver genes that may play a role in asthma, like IL-10, or antibodies to IgE and interleukins. The use of AAV5 to deliver genes to the alveolar epithelia would be of benefit in genetic diseases like alpha-1-antitrypsin. Furthermore, delivering genes to the alveolar epithelia would be of significance in several pulmonary non-genetic diseases. For example, delivering surfactant protein to premature babies or patients with ARDS; delivering genes that make antibiotic like peptides would be useful to prevent or treat pneumonia (perhaps of antibiotic-resistant organisms); delivering genes that make growth factors would be of value in emphysema; delivering genes that over-express the epithelial sodium channel or the Na-K ATPase could be used to treat cardiogenic and non-cardiogenic pulmonary edema; delivering genes that have an anti-fibrosis effect like interferon for pulmonary fibrosis would also be useful. Also, AAV5 could be used to deliver genes that may have a systemic effect like anti-hypertension drugs, insulin, coagulation factors, antibiotics, growth factors, hormones and others.

The present invention provides recombinant vectors based on AAV5. Such vectors may be useful for transducing erythroid progenitor cells or cells lacking heparin sulfate proteoglycans which is very inefficient with AAV2 based vectors. These vectors may also be useful for transducing cells with a nucleic acid of interest in order to produce cell lines that could be used to screen for agents that interact with the gene

product of the nucleic acid of interest. In addition to transduction of other cell types, transduction of erythroid cells would be useful for the treatment of cancer and genetic diseases which can be corrected by bone marrow transplants using matched donors. Some examples of this type of treatment include, but are not limited to, the introduction  
5 of a therapeutic gene such as genes encoding interferons, interleukins, tumor necrosis factors, adenosine deaminase, cellular growth factors such as lymphokines, blood coagulation factors such as factor VIII and IX, cholesterol metabolism uptake and transport protein such as EpoE and LDL receptor, and antisense sequences to inhibit viral replication of, for example, hepatitis or HIV.

10

The present invention provides a vector comprising the AAV5 virus as well as AAV5 viral particles. While AAV5 is similar to AAV2, the two viruses are found herein to be physically and genetically distinct. These differences endow AAV5 with some unique advantages which better suit it as a vector for gene therapy.

15

Furthermore, as shown herein, AAV5 capsid protein is distinct from AAV2 capsid protein and exhibits different tissue tropism. AAV2 and AAV5 likely utilize distinct cellular receptors. AAV2 and AAV5 are serologically distinct and thus, in a gene therapy application, AAV5 would allow for transduction of a patient who already  
20 possess neutralizing antibodies to AAV2 either as a result of natural immunological defense or from prior exposure to AAV2 vectors.

The present invention is more particularly described in the following examples which are intended as illustrative only since numerous modifications and variations  
25 therein will be apparent to those skilled in the art.

### EXAMPLE I

To understand the nature of AAV5 virus and to determine its usefulness as a  
30 vector for gene transfer, it was cloned and sequenced.

*Cell culture and virus propagation*

Cos and HeLa cells were maintained as monolayer cultures in D10 medium (Dulbecco's modified Eagle's medium containing 10% fetal calf serum, 100 µg/ml penicillin, 100 units/ml streptomycin and IX Fungizone as recommended by the manufacturer; (GIBCO, Gaithersburg, MD, USA) . All other cell types were grown  
5 under standard conditions which have been previously reported.

Virus was produced as previously described for AAV2 using the Beta galactosidase vector plasmid and a helper plasmid containing the AAV5 Rep and Cap  
10 genes (9). The helper plasmid was constructed in such a way to minimize any homologous sequence between the helper and vector plasmids. This step was taken to minimize the potential for wild-type (wt) particle formation by homologous recombination.

*15 DNA Cloning and Sequencing and Analysis*

In order to clone the genome of AAV5, infectious cell lysate was expanded in adherent cos cells and then suspension HeLa cells with the resulting viral particles isolated by CsCl isopycnic gradient centrifugation. DNA dot blots of Aliquots of the gradient fractions indicated that the highest concentration of viral genomes were  
20 contained in fractions with a refractive index of approx. 1.372. While the initial description of the virus did not determine the density of the particles, this value is similar to that of AAV2. Analysis of annealed virion derived DNA obtained from these fractions indicated a major species of 4.6 kb in length which upon restriction analysis gave bands similar in size to those previously reported. Additional restriction mapping  
25 indicated a unique BssHIII site at one end of the viral genome. This site was used to clone the major fragment of the viral genome. Additional overlapping clones were isolated and the sequence determined. Two distinct open reading frames (ORF) were identified. Computer analysis indicated that the left-hand ORF is approx 60% similar to that of the Rep gene of AAV2. Of the 4 other reported AAV serotypes, all have  
30 greater than 90% similarity in this ORF. The right ORF of the viral capsid proteins is also approximately 60% homologous to the Capsid ORF of AAV2. As with other

AAV serotypes reported, the divergence between AAV5 and AAV2 is clustered in multiple blocks. By using the published three dimensional structure of the canine parvovirus and computer aided sequence comparisons, a number of these divergent regions have been shown to be on the exterior of the virus and thus suggest an altered  
5 tissue tropism.

Within the p5 promoter, a number of the core transcriptional elements are conserved such as the tataa box and YY1 site around the transcriptional start site. However the YY1 site at -60 and the upstream E-Box elements are not detectable  
10 suggesting an alternative method of regulation or activation.

The inverted terminal repeats (ITRs) of the virus were cloned as a fragment from the right end of the genome. The resulting fragment was found to contain a number of sequence changes compared to AAV2. However, these changes were found  
15 to be complementary and did not affect the ability of this region to fold into a hairpin structure. Within the stem region of the hairpin two sequence elements have been found to be critical for the function of the ITRs as origins of viral replication. A repeat motif of GAGC/T which serves as the recognition site of Rep and a GGTTGAG sequence downstream of the Rep binding site which is the position of Rep's site and  
20 strand specific cleavage reaction. This sequence is not conserved between AAV5 and the other cloned AAV's suggesting that the ITRs and Rep proteins of AAV5 cannot compliment the other known AAV's.

To test the cross complementarity of AAV2 ITR containing genome and AAV5  
25 ITR containing genomes recombinant particles were packaged either using type 2 Rep and Cap or type 5 Rep and Cap expression plasmids as previously described. As shown in Fig. 2, viral particles were produced only when the respective expression plasmids were used to package the cognate ITRs. This result is distinct from that of other serotypes of AAV which have shown cross complementary in packaging.

This specificity of AAV5 Rep for AAV5 ITRs was confirmed using a terminal resolution assay which can identify the site within one ITR cleaved by the Rep protein. Incubation of the Type 5 Rep protein with a type 2 ITR did not produce any cleavage products. In contrast, addition of type 2 Rep cleaved the DNA at the expected site.

5 However AAV5 Rep did produce cleavage products when incubated with a type 5 ITR. The site mapped to a region 21 bases from the Rep binding motif that is similar to AAV2 TRS. The site in AAV2 is CGGT TGAG (SEQ ID NO: 22) but in type 5 ITR is CGGT GTGA (SEQ ID NO: 21). The ability of AAV5 Rep to cleave at a different but similarly positioned site may result in integration of AAV5 at a distinct chromosomal

10 locus compared to AAV2.

Recombinant virus produced using AAV5 Rep and Cap was obtained at a greater titer than type 2. For example, in a comparative study, virus was isolated from  $8 \times 10^7$  COS cells by CsCl banding and the distribution of the Beta galactosidase

15 genomes across the gradient were determined by DNA dot blots of aliquots of gradient fractions. DNA dot blot titers indicated that AAV5 particles were produced at a 10-50 fold higher level than AAV2.

The sequence divergence in the capsid protein ORF implies that the tissue

20 tropism of AAV2 and AAV5 would differ. To study the transduction efficiency of AAV5 and AAV2, a variety of cell lines were transduced with serial dilution's of the purified virus expressing the gene for nuclear localized Beta galactosidase activity. Approx.  $2 \times 10^4$  cells were exposed to virus in 1 ml of serum containing media for a period of 48-60 hrs. After this time the cells were fixed and stained for

25 Beta-galactosidase activity with 5-Bromo-4-chloro- 3-indolyl-b-D- galactopyranoside (Xgal) (ICN Biochemicals). Biological titers were determined by counting the number of positive cells in the different dilutions using a calibrated microscope ocular then multiplying by the area of the well. Titers were determined by the average number of cells in a minimum of 10 fields/well. Transduction of cos, HeLa, and 293, and IB3

30 cells with a similar number of particles showed approximately 10 fold decrease in titer with AAV5 compared with AAV2. In contrast MCF7 cells showed a 50-100 fold

difference in transduction efficiency. Furthermore, both vectors transduced NIH 3T3 cells relatively poorly.

A recent publication reported that heparin proteoglycans on the surface of cells are involved in viral transduction. Addition of soluble heparin has been shown to inhibit transduction by blocking viral binding. Since the transduction data suggested a difference in tissue tropism for AAV5 and AAV2, the sensitivity of AAV5 transduction to heparin was determined. At an MOI of 100, the addition of 20 $\mu$ g/ml of heparin had no effect on AAV5 transduction. In contrast this amount of heparin inhibited 90% of the AAV2 transduction. Even at an MOI of 1000, no inhibition of AAV5 transduction was detected. These data support the conclusions of the tissue tropism study, i.e. that AAV2 and AAV5 may utilize a distinct cell surface molecules and therefore the mechanism of uptake may differ as well.

AAV5 is a distinct virus within the dependovirus family based on sequence analysis, tissue tropism, and sensitivity to heparin. While elements of the P5 promoter are retained between AAV2-6 some elements are absent in AAV5 suggesting alternative mechanism of regulation. The ITR and Rep ORF are distinct from those previously identified and fail to complement the packaging of AAV2 based genomes. The ITR of AAV5 contains a different TRS compared to other serotypes of AAV which is responsible for the lack of complementation of the ITRs. This unique TRS should also result in a different integration locus for AAV5 compared to that of AAV2. Furthermore the production of recombinant AAV5 particles using standard packaging systems is approx. 10-50 fold better than AAV2. The majority of the differences in the capsid proteins lies in regions which have been proposed to be on the exterior of the surface of the parvovirus. These changes are most likely responsible for the lack of cross reactive antibodies and altered tissue tropism compared to AAV2.

From the Rep ORF of AAV2, 4 proteins are produced; The p5 promoter (SEQ ID NO: 18) produces rep 68 (a spliced site mutant) and rep78 and the p19 promoter (SEQ ID NO: 16) produces rep 40 (a spliced site mutant) and rep 52. While these



regions are not well conserved within the Rep ORF of AAV5 some splice acceptor and donor sites exist in approximately the same region as the AAV2 sites. These sites can be identified using standard computer analysis programs such as signal in the PCGENE program. Therefore the sequences of the Rep proteins can be routinely identified as in  
5 other AAV serotypes.

#### *Hemagglutination assay*

Hemagglutination activity was measured essentially as described previously (Chiorini et al 1997 J. Virol. Vol 71 6823-6833) Briefly 2 fold serial dilutions of virus  
10 in EDTA-buffered saline were mixed with an equal volume of 0.4% red blood cells in plastic U-bottom 96 well plates. The reaction was complete after a 2-h incubation at 8°C. Addition of purified AAV5 to a hemagglutination assay resulted in hemagglutination activity.

15

## **EXAMPLE II**

#### *Transduction of airway epithelial cells*

Primary airway epithelial cells were cultured and plated as previously described (Fasbender et al. J. Clin Invest. 1998 Jul 1; 102 (1): 184-93). Cells were transduced  
20 with an equivalent number of rAAV2 or rAAV5 particles containing a nuclear localized  $\beta$ -gal transgene with 50 particles of virus/cell (MOI 50) and continued in culture for 10 days.  $\beta$ -gal activity was determined following the procedure of (Chiorini et al. 1995 HGT Vol: 6 1531-1541) and the relative transduction efficiency compared. As shown in Figure 7, AAV5 transduced these cells 50- fold more efficiently than AAV2. This is  
25 the first time apical cells or cells exposed to the air have been shown to be infected by a gene therapy agent.

#### *Transduction of striated muscle*

Chicken myoblasts were cultured and plated as previously described (Rhodes &  
30 Yamada 1995 NAR Vol 23 (12) 2305-13). Cells were allowed to fuse and then transduced with a similar number of particles of rAAV2 or rAAV5 containing a nuclear

localized  $\beta$ -gal transgene as previously described above after 5 days in culture. The cells were stained for  $\beta$ -gal activity following the procedure of (Chiorini et al. 1995 HGT Vol: 6 1531-1541) and the relative transduction efficiency compared. As shown in Figure 8, AAV5 transduced these cells approximately 16 fold more efficiently than AAV2.

#### *Transduction of rat brain explants*

Primary neonatal rat brain explants were prepared as previously described (Scortegagna et al. Neurotoxicology. 1997; 18 (2): 331-9). After 7 days in culture, cells were transduced with a similar number of particles of rAAV5 containing a nuclear localized  $\beta$ -gal transgene as previously described. After 5 days in culture, the cells were stained for  $\beta$ -gal activity following the procedure of (Chiorini et al. 1995 HGT Vol: 6 1531-1541). As shown in Figure 9, transduction was detected in a variety of cell types including astrocytes, neuronal cells and glial cells.

#### *Transduction of human umbilical vein endothelial cells*

Human umbilical vein endothelial cells were cultured and plated as previously described (Gnantenko et al. J Investig Med. 1997 Feb; 45(2): 87-98). Cells were transduced with rAAV2 or rAAV5 containing a nuclear localized  $\beta$ -gal transgene with 10 particles of virus/ cell (MOI 5) in minimal media then returned to complete media. After 24 hrs in culture the cells were stained for  $\beta$ -gal activity following the procedure of Chiorini et al. (1995 HGT Vol: 6 1531-1541), and the relative transduction efficiency compared. As shown in Figure 10, AAV5 transduced these cell 5-10 fold more efficiently than AAV2.

#### *Transduction of human umbilical vein endothelial cells*

Human umbilical vein endothelial cells were cultured and plated as previously described (Gnantenko et al. J Investig Med. 1997 Feb; 45(2): 87-98). Cells were transduced with rAAV2 or rAAV5 containing a nuclear localized  $\beta$ -gal transgene with 10 particles of virus/ cell (MOI 5) in minimal media then returned to complete media. After 24 hrs in culture the cells were stained for  $\beta$ -gal activity following the procedure

of Chiorini et al. (1995 HGT Vol: 6 1531-1541), and the relative transduction efficiency compared. As shown in Figure 10, AAV5 transduced these cell 5-10 fold more efficiently than AAV2.

5

### EXAMPLE III

#### *Vector Production*

Recombinant adeno-associated viral vectors based on AAV2, AAV4, or AAV5 were prepared using high efficiency electroporation and adenovirus infection as described previously (9). All three vectors contained a nucleus-targeted *E. coli*  $\beta$ -galactosidase gene with expression driven off the Rous sarcoma virus LTR promoter (RSV). The expression cassette was flanked by AAV2 ITR sequences for rAAV2 $\beta$ gal particles and rAAV4 $\beta$ gal particles. The expression cassette was flanked by AAV5 ITR's for rAAV5 $\beta$ gal particles. The number of recombinant particles were quantified by Southern dot blot, and the biological activity was tested by X-Gal histochemical staining in a serial dilution on Cos cells. The viral titers ranged between  $2 \times 10^{11}$  to  $3 \times 10^{12}$  particles/ml and the ratio of transducing to total particles was similar to that described previously for each of the types (9,10,45). The recombinant viruses used were screened for wild-type AAV contamination by PCR, and for wild-type adenovirus by a serial dilution assay using an FITC-hexon antibody (less than  $10^3$  replication competent adenoviruses/ml) (46).

#### *Injections*

Six to 8 week old adult male C57BL/6 mice were purchased from Jackson Labs (Bar Harbor, ME) and housed at the University of Iowa Animal Care facility. All animal procedures were approved by the University of Iowa Animal Care and Use Committee. Virion injections were performed as previously described (24). Briefly, mice were anaesthetized and virions were stereotactically injected into either the right lateral ventricle or the right striatum, using a 26 gauge Hamilton syringe driven by a microinjector (Micro 1, World Precision Instruments, Sarasota, FL) at  $0.5 \mu\text{l}$  per minute. For ventricular injections,  $10 \mu\text{l}$  volumes were injected at coordinates 0.4 mm

rostral and 1.0 mm lateral to bregma, and at a 2 mm depth. For striatal injections, 5  $\mu$ l volumes were injected at coordinates 0.4 mm rostral and 2 mm lateral to bregma, and at a 3 mm depth. The doses of virion injected into the striatum, given as particle doses, were as follows: rAAV2 $\beta$ gal,  $4 \times 10^9$  (n = 5); rAAV4 $\beta$ gal,  $2 \times 10^9$  (n = 4) or  $8 \times 10^9$  (n = 3); rAAV5 $\beta$ gal,  $1.5 \times 10^{10}$  (n = 6) or  $3 \times 10^{10}$  (n = 2). For injections into the ventricle the doses were as follows: rAAV2 $\beta$ gal,  $1 \times 10^9$  (n = 3) or  $2 \times 10^9$  (n = 2); rAAV4 $\beta$ gal,  $4 \times 10^9$  (n = 8); rAAV5 $\beta$ gal,  $3 \times 10^{10}$  (n = 4). A minimum of two independent experiments was done for each virion and injection site.

#### 10 *Histochemistry*

Three or 15 weeks after injection groups of mice were perfused with 2% paraformaldehyde, the brains were removed and processed as previously described (47). 10  $\mu$ m thick coronal sections were cut at 100  $\mu$ m intervals and X-Gal histochemical staining performed to identify  $\beta$ -galactosidase expressing cells (48). For each mouse, the number of  $\beta$ -galactosidase-positive cells in every fourth section, spanning 1.3 mm of tissue rostral and 1.3 mm caudal to the injection site, were counted and summed. These sums allow quantitative comparisons among the three vectors, although they do not reflect the total number of transduced cells *in vivo*.

#### 20 *Immunofluorescent Staining*

Ten micrometer coronal cryosections of brains harvested 15 weeks after intrastriatal injection of rAAV5 $\beta$ gal were dual stained for  $\beta$ -galactosidase and either neuronal or astrocytic markers. The primary antibodies used were as follows: rabbit IgG specific for *E. Coli*  $\beta$ -galactosidase (BioDesign International, Saco MN); mouse monoclonal IgG specific for NeuN (Chemicon International, Inc., Temecula, CA), which strongly stains neuronal cell nuclei with lighter staining of the cytoplasm; and a Cy5 conjugated mouse monoclonal specific for glial fibrillary acidic protein (GFAP) (Sigma Immunocytochemicals, St. Louis, MO), an intermediate filament of astrocytes. Secondary antibodies used were ALEXA 488 goat anti-rabbit IgG (Molecular Probes, Eugene, OR) and lissamine-rhodamine goat anti-mouse IgG (Jackson ImmunoResearch Laboratories, West Grove, PA). Sections were blocked for 2h at room temperature in

phosphate-buffered saline (PBS) with 3% bovine serum albumin, 10% normal goat serum and 0.1% Triton X-100. Sections were incubated overnight with primary antibodies diluted in PBS with 3% bovine serum albumin and 0.1% Triton X-100 at 4°C, then washed and incubated with secondary antibodies in PBS with 1% normal goat serum and 0.1% Triton X-100 for 2 h at room temperature. Confocal laser microscopy was performed using 63X and 40X oil-immersion objectives on a Zeiss LSM 510 and associated software. Z-series images (0.3 to 1.0  $\mu$ m wide slices) were captured and analyzed for cellular co-localization of antigens. Colocalization of  $\beta$ -galactosidase and either NeuN or GFAP is represented in 2-color merged images from single slices within the series.

#### *Statistical Analysis*

The data was analyzed using a three-way analysis of variance with the three factors consisting of rAAV $\beta$ gal type, injection location, and time interval following virion injection. A log transformation was applied to the data to normalize the data distribution and reduce heterogeneity of group variances. Bonferroni's method was applied to each set of comparisons to adjust for the number of mice injected and to the p-values for each set of comparisons. A Bonferroni adjusted p-value < 0.05 was considered statistically significant.

#### *Quantification of Transduced Cells*

The efficiency of transduction of rAAV2, rAAV4 and rAAV5 in the brain using recombinant virion expressing the  $\beta$ -galactosidase reporter gene under control of the RSV promoter (rAAV2 $\beta$ gal, rAAV4 $\beta$ gal and rAAV5 $\beta$ gal, respectively) was evaluated. Groups of mice received either 10  $\mu$ l of vector in the right lateral ventricle, or 5  $\mu$ l of vector into the right striatum, at the particle doses stated above. Three or 15 weeks later the brains were harvested and transgene positive cells in cryosections spanning 2.6 mm rostral-caudal were quantified (Figure 11).

Three weeks after intraventricular injection, the number of rAAV4 $\beta$ gal-transduced cells was approximately 100- and 10-fold greater than for rAAV2 $\beta$ gal and

rAAV5 $\beta$ gal, respectively. rAAV2 $\beta$ gal yielded the poorest results, with the rare positive cell observed. After 15 weeks the number of rAAV5 $\beta$ gal-transduced cells was increased compared to the 3 week time point, reaching numbers similar to that of rAAV4 $\beta$ gal. This increase in rAAV5 $\beta$ gal expression nearly reached statistical  
5 significance ( $p=0.055$ ). The number of transduced cells also tended to increase from 3 to 15 weeks for rAAV2 $\beta$ gal, but remained significantly lower than for the rAAV4- and rAAV5 $\beta$ gal vectors ( $p=0.007$  and  $0.019$  respectively).

After striatal injections, strikingly greater numbers of transgene-expressing cells  
10 were detected after injection of rAAV5 $\beta$ gal compared to both rAAV2 $\beta$ gal and rAAV4 $\beta$ gal (for both,  $p<0.0001$ ). In turn, rAAV4 $\beta$ gal transduced more cells than rAAV2 $\beta$ gal by 15 weeks ( $p=0.001$ ). Comparison of the 3 and 15 week timepoints showed complete loss of rAAV2 $\beta$ gal-mediated transgene expression, but stable expression after rAAV4 $\beta$ gal injection. In contrast, there was a trend toward increased  
15 numbers of  $\beta$ -galactosidase-expressing cells from 3 to 15 weeks following rAAV5 $\beta$ gal injections.

#### *Regional Distribution of Transduced Cells*

To analyze potential regional tropisms,  $\beta$ -galactosidase-positive cells were  
20 categorized into ependyma/choroid, striatum, or other (septal area, corpus callosum, neocortex, and fornix) regions. Figure 12 illustrates the distribution of transduced cells for each virion after intraventricular or intrastriatal injections. Following intraventricular injections, transgene expressing cells were localized predominantly to the ependyma for all rAAV $\beta$ gal types at both 3 and 15 weeks (Figure 12A). Striatal  
25 injections yielded several interesting results (Figure 12B). First, rAAV2- and rAAV5 $\beta$ gal virions mediated transduction in multiple regions. Second, this data again demonstrates the global loss in rAAV2 $\beta$ gal-transduced cells in all cerebral regions from 3 to 15 weeks.

30 The patterns of transduction observed after striatal injections of the three virions are illustrated in Figure 13, which shows representative images of X-gal-stained

sections. Few blue-stained nuclei were evident in the striatum of rAAV2 $\beta$ gal -injected mice (injected dose =  $4 \times 10^9$  particles), and only at the 3 week time point (Figure 13A and B), while rAAV4 $\beta$ gal (injected dose =  $4 \times 10^9$  particles) selectively transduced the ependyma (Figure 13C). rAAV5 $\beta$ gal injections ( $1.5 \times 10^{10}$  particles) resulted in  
5 diffuse transduction in multiple cerebral regions, including the striatum (Figure 13D and E), septal region (Figure 13D) and neocortex (Figure 13F). Although the particle dose for rAAV5 $\beta$ gal was only ~4 fold greater, the relative spread of cells transduced by rAAV5 $\beta$ gal was extensive;  $\beta$ -galactosidase-expressing cells were detected 4.0 mm in the rostral-caudal, 3.5 mm dorsal-ventral, and 3.2 mm laterally, to encompass much of  
10 the injected hemisphere and portions of the medial region.

#### *Characterization of rAAV5 $\beta$ gal-transduced cells*

Previous studies have characterized the cell types transduced after parenchymal injection of rAAV2 $\beta$ gal under control of the CMV immediate early enhancer/promoter  
15 (CMVp) to be predominantly neurons, with an occasional transgene-expressing astrocyte (20,49,51). To determine which cell types were transduced by rAAV5 $\beta$ gal representative sections of brains harvested 15 weeks after intrastriatal injection were immunofluorescently stained. Confocal microscopy was performed to assess co-localization of  $\beta$ -galactosidase and representative markers. Sections were dual stained  
20 for  $\beta$ -galactosidase and either GFAP (astrocyte marker) or NeuN (neuron marker). In the striatum, many transgene-expressing cells stained positive for NeuN, indicating substantial neuronal cell transduction (Figure 14A). Transduced astrocytes were also evident in the striatum, with GFAP-positive cell processes enveloping  $\beta$ -galactosidase-positive nuclei (Figure 14B). Analyses of cells transduced in regions outside the  
25 striatum revealed that transgene-expressing cells in the cortex were also a mix of neurons and astrocytes, while those in the septal area were predominantly neurons (Figure 14C). In addition, although rAAV5 $\beta$ gal-transduced cells were noticeably more concentrated in gray matter areas, a minor proportion of transgene-positive nuclei were evident in the corpus callosum, sometimes far-removed from the injection site. In these  
30 instances, GFAP immunoreactivity identified these cells as astrocytes (Figure 14D).

In this study, CNS cell transduction with rAAV2, rAAV4 and rAAV5 virions carrying an RSV- $\beta$ -galactosidase expression cassette after intracerebral injections into the lateral ventricle or the striatum was assessed. After intraventricular injections, all three virions transduced primarily ependymal cells. Results with rAAV2 $\beta$ gal were  
5 similar to prior reports showing that transduced cells were few, and restricted to the ependyma/choroid plexus (49,50). Ependymal cell transduction was more impressive with rAAV4- and rAAV5 $\beta$ gal vectors. Since the rAAV2 $\beta$ gal and rAAV4 $\beta$ gal particles contain identical DNA sequences, differences in transduction efficiencies between these two vectors must be attributed to variations in their capsids. This implies that the  
10 rAAV4 capsid mediates more efficient entry into ependymal cells than rAAV2. rAAV5 capsid is also distinct and may likewise target ependyma more efficiently than rAAV2. Differences in the ITR region of rAAV5 $\beta$ gal may additionally influence expression. Interestingly, for rAAV5 $\beta$ gal, the number of  $\beta$ -galactosidase-positive ependymal cells increased significantly after 3 weeks, reaching levels similar to  
15 rAAV4 $\beta$ gal at 15 weeks.

Following intrastratial injections, distinct regional patterns of transduction for all three virions were observed. With rAAV4 $\beta$ gal, numerous positive cells lined the ventricles, with very few transgene-expressing cells in the parenchyma. In contrast,  
20 rAAV2 $\beta$ gal and rAAV5 $\beta$ gal vectors transduced predominantly parenchymal cells, and unlike rAAV2 $\beta$ gal (20,49,51), rAAV5 $\beta$ gal transduced a significant proportion of astrocytes as well as neurons. Moreover, rAAV5 $\beta$ gal transduced a greater number of cells, over a larger volume of tissue compared to rAAV2 $\beta$ gal.

25 When compared to AAV2, heterogeneities in the capsid-encoding regions, heparin-insensitive transduction, and differential abilities to transduce cell lines *in vitro* together strongly implicate different receptor requirements for cell entry by AAV5 (10,45,56). The enhanced parenchymal cell transduction observed for rAAV5 $\beta$ gal compared to rAAV2 $\beta$ gal, as well as the diffuse and widespread pattern of transduction  
30 mediated by rAAV5 $\beta$ gal may similarly reflect distinct receptor requirements. In the rat brain, rAAV2 $\beta$ gal particles have been shown to preferentially bind neurons, but not



glial cells, within minutes of injection, and be transported within 30 minutes to neuronal cell nuclei (57). Considering that neuronal subtypes in adult rodent brain express integral membrane heparan sulfate proteoglycans such as syndecans (58,59) and glypican-1 (60), it is conceivable that AAV2 binds strongly to and enters neurons surrounding the injection site. Moreover AAV2 particles may become sequestered in extracellular HSPG in a way that limits vector diffusion and reduces transduction efficiency. An ability of rAAV5 $\beta$ gal to travel in a less restricted fashion may explain these observations of widespread transduction in comparison to rAAV2 $\beta$ gal.

Following intrastriatal injection loss of rAAV2 $\beta$ gal-transduced cells over time consistent with observations of others (20,49,50,51,62,63) was observed. In contrast, transgene expression after rAAV5 $\beta$ gal injection was stable over the time-course of our study. rAAV5 could target to cell subsets better able to sustain RSVp-driven transcription, or there could be positive influences of the AAV5 ITRS on either genome stability or RSV promoter activity.

These experiments explored the use of rAAV5 as a vector for gene transfer to the CNS. rAAV5 $\beta$ gal transduced large numbers of cells, with lasting expression in both neuronal and glial types. More importantly, rAAV5 $\beta$ gal exhibited an extensive transduction volume. Vector diffusion is an extremely valuable feature for gene therapy of CNS diseases exhibiting widespread pathology, such as the neurodegenerative aspect of the lysosomal storage diseases. This characteristic coupled to persistent expression could reduce the need for multiple injection sites and repeat injections, and their obvious associated risks.

#### EXAMPLE IV

##### *Human airway epithelia*

Airway epithelial cells were obtained from surgical polypectomies of non-CF patients or from trachea and bronchi of lungs removed for organ donation. Cells were isolated by enzyme digestion as previously described (67). Freshly isolated cells were

seeded at a density of  $5 \times 10^5$  cells/cm<sup>2</sup> onto collagen-coated, 0.6 cm<sup>2</sup> diameter Millicell polycarbonate filters (Millipore Corp., Bedford, MA). The cells were maintained at 37 °C in a humidified atmosphere of 7% CO<sub>2</sub> and air. Twenty-four hours after plating, the mucosal media was removed and the cells were allowed to grow at the air-liquid interface (68, 69). The culture medium consisted of a 1:1 mix of DMEM/Ham's F12, 5% Ultrosor G (Biosepra SA, Cedex, France), 100 U/ml penicillin, 100 µg/ml streptomycin, 1% nonessential amino acids, and 0.12 U/ml insulin. Airway epithelia were allowed to reach confluence and develop a transepithelial electrical resistance (Rt), indicating the development of tight junctions and an intact barrier. Epithelia were allowed to differentiate by culturing for at least 14 days after seeding and the presence of a ciliated surface was tested by scanning electron microscopy (70).

#### *Recombinant adeno-associated viruses*

Recombinant AAV vectors expressing β-galactosidase, AAV2/βGal, AAV4/βGal, AAV5/βGal, were prepared using high efficiency electroporation and packaging initiated by adenovirus infection and characterized as described previously (45). Briefly, rAAV particles were produced by electroporating  $1 \times 10^8$  exponentially growing Cos cells with 400 µg of a 1:1 mixture of pAAV2RnLacZ and pSV40oriAAV2 for production of AAV2, pAAV2RnLacZ and pSV40oriAAV4 for AAV4, or pAAV5RnLacZ and pSV40oriAAV5 for AAV5 in 1X RPMI (2.5 ml of 2X RPMI, 1 ml FCS, 1.5 ml H<sub>2</sub>O, and 50 µl of 1M Hepes pH 7.4) and incubated on ice for 10 min prior to electroporation. Electroporation was performed in a 4 mm gap cuvette (BioRad Richmond CA) containing 0.5 mls of the cell DNA mixture using a BTX 600 electroporator. Conditions used for electroporation were 300 Volt, 2100 µF, 48 ohms. Following electroporation the cells were incubated on ice for 10 min then plated into ten 15 cm dishes. The following day the medium was replaced and the cells allowed to recover. Approximately 30-50% of the cells which were initially electroporated reattached to the plates and 90% of these cells show strong expression of the β-gal reporter gene. Two days later, the plates were infected with approximately  $5 \times 10^9$  PFU (MOI of 10) of wild-type adenovirus type 5 for 1 h. in serum free media and then supplemented with D10 media. Seventy-two hours post infection the cells were

harvested by scraping and the virus and the cells pelleted by low speed centrifugation. The pellet is resuspended in 7.5 ml of TD buffer for every 10 plates (TD = 140 mM NaCl, 5 mM KCl, 0.7 mM K<sub>2</sub>HPO<sub>4</sub>, 25 mM Tris/HCl, pH 7.4). Trypsin (0.5 volumes of 0.25%) and sodium deoxycholate (0.5 volumes of 10%) are added to the suspension, 5 gently mixed and incubated at 37 °C for 30 min. The lysate is then homogenized thoroughly (approximately 20 strokes in a Wheaton B homogenizer). CsCl is next added to a final density of 1.4 g/cm<sup>3</sup> and the homogenate is distributed into two polyallomer tubes and centrifuged in a SW40.1 swinging bucket rotor at 38,000 RPM for 65 hr at 20 °C. The pellicle at the top of the gradient is removed using a pasture 10 pipette and the gradients fractionated by side puncture. Fractions with a refractive index of 1.373-1.371 were pooled for AAV2 and AAV5 and 1.378-1.376 for AAV4, and centrifuged again using an SW50.1 rotor, and fractionated as described above. Refractive indices were determined using a Zeiss refractometer.

15 Recombinant viruses were titered by Southern blot, and X-Gal staining in a serial dilution on COS-7 cells tested their biological activity. The viral titers ranged between  $4 \times 10^{12}$  and  $8 \times 10^{12}$  particles/ml. The particle to transduction unit ratio on these cells was similar to that previously reported for all 3 viruses on Cos cells (about  $10^4$  to 1). The recombinant viruses used were screened for wild-type AAV 20 contamination by PCR, and for wild-type adenovirus by a serial dilution assay using a FITC-hexon antibody (less than  $10^3$  replication competent adenoviruses/ml) (70).

#### *Viral infection and binding assays*

Five hundred virions of the recombinant AAV/per cell (in phosphate-buffered 25 saline) were added to the apical surface. Following the indicated incubation time, the viral suspension was removed and the epithelia were rinsed twice with PBS. After infection, the epithelia were incubated at 37 °C for an additional fourteen days.

To assess binding to airway epithelia, the epithelia were incubated for 30 min at 30 4 °C with 500 virions/cell of AAV2/βGal, AAV4/βGal or AAV5/βGal. The epithelia were then rinsed, and cell-associated AAV viral DNA was measured from cell lysates

of seven epithelia per dot. Samples were subjected to 3 freeze/thaw cycles and then blotted onto a nylon membrane (Ambion, Austin, TX). Detection of the AAV viral DNA was done by hybridizing with a  $^{32}\text{P}$ -labeled pCMV $\beta$ gal. Unhybridized probe was washed as follows: two washes with 2% SSC and 0.1X SDS at room temperature  
5 for 15 min, one wash with 0.5X SSC and 0.1% SDS at 55 °C for 1 hr, and finally one wash with 0.5X SSC and 0.1% SDS at 65 °C for 30 min. Dot blots were developed and quantitated using a PhosphorImager (Molecular Dynamics, Sunnyvale, CA) (71).

#### *Measurement of $\beta$ -galactosidase activity*

10 Total  $\beta$ -galactosidase activity was measure using a commercially available method (Galacto-Light, Tropix, Inc., Bedford, MA). Briefly, after rinsing with PBS, cells were removed from filters by incubation with 120  $\mu\text{l}$  lysis buffer (25 mM Tris-phosphate, pH 7.8; 2 mM DTT; 2 mM 1, 2-diaminocyclohexane-N,N,N',N'-tetraacetic acid; 10% glycerol; and 1% Triton X-100) for 15 min. Light emission was quantified  
15 in a luminometer (Analytical Luminescence Laboratory, San Diego CA). To histochemically detect  $\beta$ -galactosidase activity, the chromogenic reagent X-Gal (5-bromo-4-chloro-3-indonyl- $\beta$ -D-galactopyranoside, Boehringer Mannheim) was used. Human airway epithelia and murine lungs were fixed with 1.8% formaldehyde and 2% glutaraldehyde, and then incubated for 16 hr at 37° C with 313  $\mu\text{l}$  of 40 mg/ml X-Gal in  
20 DMSO dissolved in 12.5 ml of PBS (pH 7.8).

#### *Studies in mice*

For *in vivo* analysis, 6-8 week old C57BL/6 mice (The Jackson Laboratory, Bar Harbor, ME) were studied. Mice were lightly anesthetized using a methoxyflurane  
25 chamber. Recombinant AAV2 and AAV5 ( $1 \times 10^{10}$  particles) were administered intranasally in two 62.5  $\mu\text{l}$  instillations delivered 5 min apart. The experiment was performed with five animals per group. Twenty-eight days after vector administration, animals were sacrificed with  $\text{CO}_2$ . PBS (10 ml) was instilled into the right ventricle and then the lungs and heart were removed intact. The trachea was intubated and  
30 instilled at 10 cm of pressure with the following solutions in order: PBS, 4% paraformaldehyde, PBS, and stained overnight with X-Gal stain and finally rinsed with

PBS. Lungs were cryosectioned and sections were analyzed by two independent reviewers that were unaware of the experimental identity of the samples. The reviewers counted the number of blue nuclei of  $\beta$ gal-expressing cells from a 5  $\mu$ m slice obtained every 50  $\mu$ m ( $n=20$  fields/lung). The total number of airway epithelial cells was  
5 estimated by dividing the surface of the epithelia ( $\pi 2r$ ) by (4.9  $\mu$ m), an estimate of the diameter of the airway epithelial cells ( $2425.3 \pm 20$  airway cells/field).

AAV5 can mediate gene transfer through the apical surface of human airway epithelia. Because AAV2, AAV4, and AAV5 have different tropism in cell lines, the  
10 efficiency of these different serotypes on primary cultures of differentiated human airway epithelia was compared. Epithelia were transduced for 12 hours at a relatively low particle per cell ratio (500 particles/cell) with an estimated MOI of less than 1. To allow for maximal expression, the epithelia were studied 2 weeks after infection. Quantification of the  $\beta$ -galactosidase activity showed that AAV5-transduced cells  
15 generated approximately 50 fold greater activity than AAV2 or AAV4 transduced cells (Fig 15E). To histochemically detect the  $\beta$ -galactosidase activity, we stained the epithelia with a chromogenic reagent X-gal. Similar to the quantitative analysis, figure 15B & 15C shows only minimal gene transfer in epithelia transduced by AAV2/ $\beta$ Gal or AAV4/ $\beta$ Gal compared to epithelia transduced with AAV5/ $\beta$ Gal (Fig 15D). To rule out  
20 the possibility of pseudo-transduction by protein transfer, the epithelia was assayed 1 hour after the application of the AAV vectors, no  $\beta$ -gal activity was detected over background.

*AAV5 binds to the apical surface of well-differentiated human airway epithelia*

25 The hypothesis that the improved transduction efficiency of AAV5/ $\beta$ Gal relied on increased binding to well-differentiated airway epithelia was tested. Epithelia were incubated for 30 min with 500 particles per cell of AAV2/ $\beta$ Gal, AAV4/ $\beta$ Gal or AAV5/ $\beta$ Gal, then rinsed. Cell-associated AAV was estimated by dot blot analysis. Figure 16 shows that differentiated airway epithelia bound AAV5 derived vector  
30 approximately seven-fold better than AAV2/ $\beta$ Gal. Of interest, AAV4/ $\beta$ Gal also bound to the apical surface five times more efficiently than AAV2/ $\beta$ Gal. These data may

explain some of the advantage of AAV5 over AAV2-derived vectors in mediating gene transfer to the airway epithelia.

5 *Effect of dose and incubation time on AAV5 infection of the apical surface of human airway epithelia*

Since AAV5 appeared to bind and mediate gene transfer to the airway epithelia more efficiently than AAV2, the effect of dose of the virus was examined. Figure 17 shows that in a range of 0.5 to 5000 particles/cell, AAV5 always outperformed AAV2/ $\beta$ Gal. The course of AAV5-mediated expression of  $\beta$ -galactosidase *in vitro* over a month period was also tested. The level of  $\beta$ -galactosidase expression was stable over 28 days ( $3.4 \times 10^7 \pm 1.4 \times 10^7$  L.U./mg and  $3.18 \times 10^7 \pm 1.1 \times 10^7$  L.U./mg for 10 and 28 days respectively).

The effect of incubation time for the virion (AAV5/ $\beta$ Gal) on airway epithelia was tested. Figure 18 shows that contrary to what is seen with recombinant adenovirus and AAV2, incubation of airway epithelia with recombinant AAV5 resulted in similar levels of gene transfer with short incubation, 30 min or a prolonged incubation, 12 h. This is in agreement with the increased affinity found for AAV5 compared to AAV2 and adenoviruses and more importantly it suggest there may be an apical receptor for AAV5.

*AAV5 infection of the apical surface of human airway epithelia is not sensitive to heparin competition.*

The low level transduction of airway epithelia by AAV2/ $\beta$ Gal is thought to be the result of poor virus binding because the apical membrane of airway epithelia expresses very low levels of HSP and  $\alpha$ V $\beta$  integrins that may mediate AAV2 binding (54,74). To test the effect of heparin competition on AAV2/ $\beta$ Gal and AAV5/ $\beta$ Gal transduction of human airway epithelia, the viruses were pre-incubated with 20  $\mu$ g/ml of soluble heparin. Competition with soluble heparin had minimal effect on the already low level of AAV2/ $\beta$ Gal-mediated gene transfer suggesting that the observed low level transduction was not receptor-mediated (Fig 19A). However more importantly, heparin

competition did not inhibit AAV5/ $\beta$ Gal-mediated gene transfer to airway epithelia. These data show a novel receptor-mediated pathway for AAV5 binding and infection via the apical surface of human airway epithelia.

5 *AAV 5 mediates gene transfer through the basolateral surface in a heparin sulfate independent manner*

The binding of AAV5 to the apical membrane suggests a novel receptor. To test if the receptor for AAV5 is present on the basolateral surface, the transduction experiments were repeated as described in the previous section but vector was applied  
10 from the basolateral side. Because AAV2 can infect via the basolateral side, this experimental design also allowed investigation of whether or not AAV5 had an advantage over AAV2 once they were both in the cell. Briefly, the epithelia were turned upside down and 500 virions/cell of AAV5/ $\beta$ Gal or AAV2/ $\beta$ Gal was carefully applied in a volume of 25  $\mu$ l to the bottom of the Millipore filter. After 30 min, the  
15 epithelia were rinsed thoroughly. To allow for maximal expression, the epithelia were studied 2 weeks after infection. Figure 19B shows that similar levels of  $\beta$ -galactosidase activity were detected in airway epithelia transduced with either AAV2/ $\beta$ Gal or AAV5/ $\beta$ Gal. These data suggest that both viruses work equally well when applied to the basolateral side. To test the mechanism of uptake, the studies were  
20 repeated in the presence of soluble heparin. As previously reported, basolateral infection of the airway epithelia by AAV2 was competed off by soluble heparin (74). However, the AAV5/ $\beta$ Gal transduction via the basolateral surface was not blocked by heparin competition. These data show that that AAV5 binds to a different receptor than AAV2 that is present both on the apical and basolateral surfaces of human airway  
25 epithelia.

*AAV 5 mediated gene transfer to the airways in vivo*

These data demonstrate improved gene transfer of human ciliated airway epithelia with AAV5 compared to AAV2. To compare the transduction efficiency of  
30 AAV5 and AAV2 *in vivo*, administered either AAV2 or AAV5 ( $1 \times 10^{10}$  particles) was administered to 6-8 week old C57BL/6 mice, in a total volume of 125  $\mu$ l via nasal

instillation. After 30 days the mice were sacrificed, and the lungs were fixed and stained with X-Gal as previously described (71). A relatively low viral input was chosen to maximize the difference between specific receptor binding and non-specific binding that may occur when the viral concentrations are very high (70,73,74,77). Only minimal transduction in mice treated with AAV2/ $\beta$ Gal was observed (Fig 20). In contrast, a significant increase in the number of blue cells in the lungs of mice treated with AAV5 was observed. A 15 fold increase over AAV2 transduction was observed when alveolar cells were transduced with AAV5. These data confirm the *in vitro* observation that AAV5 is more efficient at mediating gene transfer to the luminal surface of airway epithelia than AAV2 and suggest that murine airway epithelia express the receptor for AAV5.

The data presented in this Example suggest that the capsid of AAV5 is sufficiently different from that of AAV2 to allow for efficient binding and infection of human airway epithelia. While previous research has demonstrated transduction of airway epithelial cells with AAV2 those studies have required either very high MOI's and/or prolonged incubation times. The present invention shows that human and murine airway epithelia can be more efficiently transduced by AAV5. Furthermore the data suggest a novel receptor present both in the apical and basolateral surface of airway epithelia.

## EXAMPLE V

### *Preparation of viral vectors*

AAV5 expressing nuclear targeted  $\beta$ -galactosidase driven off of a Rous sarcoma virus promoter was prepared. Virus was concentrated and suspended in 3% sucrose in phosphate buffered saline prior to *in vivo* use. AAV5 titres were approximately  $1 \times 10^8$  infectious units/ml as assessed by  $\beta$ -galactosidase histochemistry of COS cells transfected with serial dilutions of the viruses. For most of the experiments, the neuronal tracer cholera toxin subunit b (CTb) was added to the viral suspension at a concentration of 1  $\mu$ g/ $\mu$ l so that CTb immunoreactivity could be used to



independently visualize cerebellar injection sites and distinguish transport and spread of virus outside of the injection site from transduction within the primary injection site itself. CTb is the nontoxic, nonbiologically active subunit of cholera toxin and is presumed to be inert in neuronal tracing experiments. Nonetheless, to insure that CTb had no effects on results AAV5, several animals were injected with virus alone (no CTb).

#### *Cerebellar Injections and Preparation of Tissue*

Young adult C57Bl6 mice weighing 20-25 g were anesthetized with ketamine/xylazine, a burr hole was drilled at the midline posterior occipital bone overlying the cerebellar anterior lobe and pressure injections totaling 2 $\mu$ l were made into a single cerebellar lobule using a Hamilton syringe cemented with a glass micropipette tip. After survival periods, 7 weeks for AAV5, animals were reanesthetized and transcardially perfused with cold phosphate buffered saline followed by 4% paraformaldehyde in 0.1 M phosphate buffer, pH 7.4. Cerebella, brainstems and thoracolumbar spinal cords were removed and postfixed in the 4% paraformaldehyde overnight at 4 °C, cryoprotected for 1-3 days in 30% sucrose in phosphate buffered saline at 4 °C and then sectioned on a cryostat at 50  $\mu$ m thickness (cerebellum/brainstem sagittally and spinal cord longitudinally).

#### *Histochemistry and immunofluorescence*

Gene transfer was determined by processing every other section for  $\beta$ -galactosidase staining with 5-bromo-4-chloro-4-indolyl  $\beta$ -D-galactoside (X-Gal) according to Terashima 1997. Transport and spread of virus was then determined by comparing the X-gal processed sections to adjacent sections that had been processed for CTb immunohistochemistry according to Alisky and Tolbert (1994). Finally, neuronal versus glial gene transduction was determined by dual staining immunofluorescence for  $\beta$ -galactosidase and neuronal and glial markers on selected cerebellar sections. Glial fibrillary acid protein (GFAP) was used as the glial marker and calbindin was used as the neuronal marker; colocalization was then determined by confocal microscopy.

*Injection sites*

As determined by CTb immunoreactivity, all injection sites were approximately the area of a single cerebellar lobule in the anterior lobe (lobules II, III, IV or V) encompassing most of the anteroposterior and mediolateral extent of the lobule within the vermis. In some cases, injections encompassed the dorsal half of one lobule and the ventral half of another lobule for a net injection of a single lobule. Injections filled the molecular layer, Purkinje cell layer, granule cell layer and white matter of the arbor vitae but did not extend to the deep cerebellar nuclei. Outside the injection site, the CTb retrogradely labeled precerebellar neurons in the cuneate, vestibular, olivary, reticular and spinal nuclei, thus mapping an extensive pool of neurons which could be potentially transfected via retrograde axonal transport of virus.

*AAV5 $\beta$ gal Transduction*

There were 12 mice in the AAV arm, seven with AAV5-CTb and five with AAV5 alone, all sacrificed at 7 weeks postinjection. Neuronal cell types could be unequivocally identified because of the unique morphology and position of different neuronal classes within the cerebellar cortex. Stellate neurons are always outermost in the molecular layer, basket cells are in the inner part of the molecular layer, Purkinje cells are always a monolayer and Golgi and granule neurons are exclusively in the granule cell layer. There was no difference between AAV5-CTb and AAV5 alone and tropism of the two vectors was similar. There was extensive AAV5- $\beta$ gal gene transfer to Purkinje cells, stellate and basket cells and Golgi neurons but only minimal transduction of granule cell neurons (Figure 21). By immunofluorescence, transduction was exclusively neuronal (Figure 22 and 23)

25

AAV5 showed little retrograde transport but much greater physical spread. The only retrograde transport was to deep cerebellar nuclei; the brainstem vestibular nuclei transduced by the FTV were not transduced by AAV5. However, AAV5- $\beta$ galactosidase expressing Purkinje cells could be seen in several lobules far beyond the single lobule injection sites, sometimes the entire anterior and posterior lobes. Moreover AAV5- $\beta$ gal expressing cells could be seen in the overlying inferior colliculi in x of the

30

animals, clearly diffusion rather than retrograde axonal transport as there are no axon projections from the inferior colliculus into the cerebellum.

Throughout this application, various publications are referenced. The  
5 disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

Although the present process has been described with reference to specific  
10 details of certain embodiments thereof, it is not intended that such details should be regarded as limitations upon the scope of the invention except as and to the extent that they are included in the accompanying claims.

#### References:

1. Arella M., Garzon S., Bergeron J., and Tijssen P. Handbook of Parvoviruses. Vol. 1. ed. P. Tijssen. Boca Raton, Florida, CRC Press, 1990.
2. Bachmann PA, Hoggan MD, Kurstak E., Melnick JL, Pereira HG, Tattersall P., and Vago C. 1979. Intervirology 11: 248-254.
3. Bantel-Schaal U. and Stohr M. 1992. J. Virol. 66: 773-779.
4. Chang LS, Shi Y., and Shenk T. 1989. J. Virol. 63: 3479-88.
5. Chejanovsky N. and Carter BJ 1989. Virology 173: 120-128.
6. Chejanovsky N. and Carter BJ 1989. Virology 171: 239-247.
7. Chiorini JA, Wiener SM, Kotin RM, Owens RA, Kyöstiö SRM, and Safer B. 1994. J. Virol. 68: 7448-7457.
8. Chiorini JA, Weitzman MD, Owens RA, Urcelay E., Safer B., and Kotin RM 1994. J. Virol. 68: 797-804.
9. Chiorini JA, Wendtner CM, Urcelay E., Safer B., Hallek M., and Kotin R.M. 1995. Human Gene Therapy 6: 1531-1541.
10. Chiorini JA, Yang L., Safer B., and Kotin RM 1995. J. Virol. 69: 7334-7338.

11. Dixit M., Webb MS, Smart WC, and Ohi S. 1991. *Gene* **104**: 253-7.
12. Fisher RE and Mayor HD 1991. *J Theor Biol* **149**: 429-39.
13. Flotte TR, Afione SA, Conrad C., McGrath SA, Solow R., Oka H., Zeitlin PL, Guggino WB, and Carter BJ 1993. *Proc. Natl. Acad. Sci.* **90**: 10613-10617.
14. Flotte, T.R., S.A. Afione, R. Solow, M.L. Drumm, D. Markakis, W.B. Guggino, P.L. Zeitlin, and B.J. Carter. 1993. *J Biol Chem* **268**: 3781-90.
15. Hermonat, P.L., M.A. Labow, R. Wright, K.I. Berns, and N. Muzyczka. 1984. *J. Virol.* **51**: 329-339.
16. Hermonat, P.L. and N. Muzyczka. 1984. *Proc Natl Acad Sci USA* **81**: 6466-70.
17. Hunter, L.A. and R.J. Samulski. 1992. *J. Virol.* **66**: 317-24.
18. Ito, M. and H.D. Mayor. 1968. *J. Immuno.* **100**: 61-68.
19. Janik, J.E., M.M. Huston, K. Cho, and J.A. Rose. 1989. *Virology* **168**: 320-9.
20. Kaplitt, M.G., P. Leone, R.J. Samulski, X. Xiao, D.W. Pfaff, K.L. O'Malley, and J.M. Duing. 1994. *Nature Genetics* **8**: 148-154.
21. Kotin, R.M., M. Siniscalco, R.J. Samulski, X. Zhu, L. Hunter, C.A. Laughlin, S. McLaughlin, N. Muzyczka, M. Rocchi, and K.I. Berns. 1990. *Proc. Natl. Acad. Sci. (USA)* **87**: 2211-2215.
22. Laughlin, C.A., N. Jones, and B.J. Carter. 1982. *J. Virol.* **41**: 868-76.
23. Laughlin, C.A., M.W. Myers, D.L. Risin, B.J. Carter. 1979. *Virology* **94**: 162-74.
24. McCarty, D.M., J. Pereira, I. Zolotukhin, X. Zhou, J.H. Ryan, and N. Muzyczka. 1994. *J. Virol.* **68**: 4988-4997.
25. Mendelson, E., J.P. Trempe, and B.J. Carter. 1986. *J. Virol.* **60**: 823-832.
26. Mizukami, H., N.S. Young, and K.E. Brown. 1996. *Virology* **217**: 124-130.
27. Muster, C.J., Y.S. Lee, J.E. Newbold, and J. Leis. 1980. *J. Virol.* **35**: 653-61.
28. Muzyczka, N. 1992. *Curr Top Microbiol Immunol* **158**: 97-129.
29. Parks, W.P., J.L. Melnick, R. Rongey, and H.D. Mayor. 1967. *J. Virol.* **1**:

- 171-180.
30. **Podsakoff, G., K.K. Jr Wong, and S. Chatterjee.** 1994. *J. Virol.* **68**: 5656-5666.
  31. **Rose, J.A., M.D. Hoggan, F. Koczot, and A.J. Shatkin.** 1968. *J. Virol.* **2**: 999-1005.
  32. **Russell, D.W., A.D. Miller, and I.E. Alexander.** 1994. *Proc. Natl. Acad. Sci. USA* **91**: 8915-8919.
  33. **Ryan, J.H., S. Zolotukhin, and N. Muzyczka.** 1996. *J. Virol.* **70**: 1542-1553.
  34. **Samulski, R.J., K.I. Berns, M. Tan, and N. Muzyczka.** 1982. *Proc Natl Acad Sci USA* **79**: 2077-81.
  35. **Samulski, R.J., L.S. Chang, and T. Shenk.** 1989. *J. Virol.* **63**: 3822-8.
  36. **Sanes, J.R., J.L.R. Rubenstein, and J.F. Nicocas.** 1986. *EMBO* **5**: 3133-3142.
  37. **Senapathy, P., J.D. Tratschin, and B.J. Carter.** 1984. *J Mol Biol* **179**: 1-20.
  38. **Tratschin, J.D., I.L. Miller, and B.J. Carter.** 1984. *J. Virol.* **51**: 611-619.
  39. **Trempe, J.P. and B.J. Carter.** 1988. *J. Virol.* **62**: 68-74.
  40. **Trempe, J.P., E. Mendelson, and B.J. Carter.** 1987. *Virology* **161**: 18-28.
  41. **Walsh, C.E., J.M. Liu, X. Xiao, N.S. Young, A.W. Nienhuis, and R.J. Samulski.** 1992. *Proc Natl Acad Sci USA* **89**: 7257-61.
  42. **Winocour, E., M.F. Callahan, and E. Huberman.** 1988. *Virology* **167**: 393-9.
  43. **Jaksch, M., K.D. Gerbitz, and C. Kilger.** 1995. *Clin. Biochem.* **28**:503-509
  44. **Burcin, M.M., O'Malley, B.W. and S.Y. Tsai.** 1998. *Frontiers in Bioscience* **3**:1-7.
  45. **Chiorini J.A., Kim F., Yang L., Kotin R.M.** 1999 *J Virol* **73**(2):1309-1319.
  46. **Hehir K.M., Armentano D., Cardoza L.M., Choquette T.L., Berthelette P.B., White G.A., Couture L.A., Everton M.B., Keegan J., Martin J.M., Pratt D.A., Smith M.P., Smith A.E., Wadsworth S.C.** 1996 *J Virol* **70**(12):8459-8467.
  47. **Ghodsi A., Stein, C., Derksen T., Yang, G., Anderson R.D., Davidson B.L.**

- 1998 Hum Gene Ther 9:2331-2340.
48. Davidson BL, Doran SE, Shewach DS, Latta JM, Hartman JW, Roessler BJ. 1994 Exp Neurol 125:258-267.
  49. McCown TJ, Xiao X, Li J, Breese GR, Samulski RJ. 1996 Brain Res 713:99-107.
  50. Lo WD, Qu G, Sferra TJ, Clark R, Chen R, Johnson PR. 1999 Hum Gene Ther 10:201-213.
  51. Mandel RJ, Rendahl KG, Spratt SK, Snyder RO, Cohen LK, Leff SE. 1998 J Neurosci 18(11):4271-4284.
  52. Summerford C, Samulski RJ. 1998 J Virol 72(2):1438-1445.
  53. Qing K, Mah C, Hansen J, Zhou S, Dwarki V, Srivastava A. 1999 Nat Med 5(1):71-77.
  54. Summerford C, Bartlett JS, Samulski RJ. 1999 Nat Med 5(1):78-82.
  55. Qiu J, Brown KE. 1999 Virology 264(2):436-440.
  56. Bantel-Schaal U, Delius H, Schmidt R, zur Hausen H. 1999 J Virol 73(2):939-947.
  57. Bartlett JS, Samulski RJ, McCown TJ. 1998 Hum Gene Ther 9(8):1181-1186.
  58. Hsueh Y-P, Yang F-C, Kharazia V, Naisbitt S, Cohen AR, Weinberg RJ, Sheng M 1998 J Cell Biol 142(1):139-151.
  59. Hsueh Y-P, Sheng M. 1999 J Neurosci 19(17):7415-7425.
  60. Liang Y, Annan RS, Carr SA, Popp S, Mevissen M, Margolis RK, Margolis RU. 1999 J Biol Chem 274(25):17885-17892.
  61. Fan D-S, Ogawa M, Fujimoto K-I, Ikeguchi K, Ogasawara Y, Urabe M, Nishizawa M, Nakano I, Yoshida M, Nagatsu I, Ichinose H, Nagatsu T, Kurtzman GJ, Ozawa K 1998 Hum Gene Ther 9:2527-2535.
  62. Doll RF, Crandall JE, Dyer CA, Aucoin JM, Smith FI. 1996 Gene Ther 3:437-447.
  63. Klein RL, Meyer EM, Peel AL, Zolotukhin S, Meyers C, Muzyczka N, King MA. 1998 Exp Neurol 150:183-194.
  64. Bajocchi G, Feldman SH, Crystal RG, Mastrangeli A. 1993 Nat Genet

3:229-234.

65. **Ghodsi A., Stein C., Derksen T., Martins I., Anderson RD, & Davidson BL** 1999 *Exp Neurol* 160, 109-116.
66. **Johansson CB, Momma S, Clarke DL, Risling M, Lendahl U, Frisen J** 1999 *Cell* 96(1):25-34.
67. **Rich DP, Couture LA, Cardoza LM, Guiggio LM, Armentano D., Espino PC, Hehir K., Welsh MJ, Smith AE, and Gregory RJ** 1993 *Hum.Gene Ther.* 4:461-476.
68. **Kondo M., Finkbeiner WE, and Widdicombe JH.** 1991. *Am.J.Physiol.* 261:L106-L117
69. **Yamaya, M., Finkbeiner WE, Chun SY, and Widdicombe JH** 1992 *Am.J.Physiol.* 262:L713-L724
70. **Zabner, J., Zeiher BG, Friedman E, and Welsh MJ** 1996 *J.Virol.* 70:6994-7003.
71. **Walters, RW, Duan D., Engelhardt JF, and Welsh MJ.** 1999. *J. Virol.* (In Press)
72. **Bergelson, JM, Cunningham JA, Droguett G., Kurt-Jones EA, Krithivas A., Hong JS, Horwitz MS, Crowell RL, and Finberg RW** 1997 *Science* 275:1320-1323.
73. **Teramoto, S., Bartlett JS, McCarty DXX, Samulski RJ, and Boucher RC** 1998 *J Virol* 72:8904-8912.
74. **Duan, D., Yue Y., Yan Z., McCray PBJr, and Engelhardt JF.** 1998 *Hum Gene Ther* 9:2761-2776.
75. **Walters, RW, Grunst T., Bergelson JM, Finberg RW, Welsh MJ, and Zabner J.** 1999 *J. Biol. Chem.* 274:10219-10226.
76. **Wang G., Davidson BL, Melchert P., Slepishkin VA, van Es HH, Bodner M., JollyDJ, and McCray PBJr.** 1998. *Journal of Virology* 72:9818-9826.
77. **Halbert CL, Standaert TA, Aitken ML, Alexander IE, Russell DW, and Miller AD** 1997. *J.Virol.* 71:5932-5941.
78. **Fisher, KJ, Jooss K., Alston J., Yang Y., Haecker SE, High K., Pathak R., Raper SE, and Wilson JM** 1997 *Nat Med* 3:306-312.

79. **Guy J., Qi X., Muzyczka N., and Hauswirth WW** 1999 Arch Ophthalmol 117:929-937.
80. **Snyder RO, Miao CH, Patijn GA, Spratt SK, Danos O., Nagy D., Gown AM, Winther B., Meuse L., Cohen LK, Thompson AR, and Kay MA** 1997. Nat.Genet. 16:270-276.
81. **Girod A., Ried M., Wobus C., Lahm H., Leike K., Kleinschmidt J., Deleage G., and Hallek M.** 1999 Nat Med 5:1052-1056.
82. **Bartlett JS, Kleinschmidt J., Boucher RC, and Samulski RJ** 1999 Nat Biotechnol 17:181-186.
83. **Xiao W., Chirmule N., Berta SC, McCullough B., Gao G., and Wilson JM** 1999 J Virol 73:3994-4003.
84. **Rutledge EA, Halbert CL, and Russell DW** 1998 J Virol 72:309-319.
85. **Xie Q. and Chapman MS** 1996 J Mol Biol 264:497-520.
86. **Alisky J.M. and Tolbert D.M.** 1994 Journal of Neuroscience Methods 52: 143-148.



What is claimed is:

1. A method of delivering a nucleic acid to a cell, comprising administering to the cell an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to the cell.
2. A method of delivering a nucleic acid to a cell in a subject comprising administering to the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to a cell in the subject.
3. The method of claim 1, wherein the cell is selected from the group consisting of: alveolar cell, cerebellar cell, and ependymal cell.
4. The method of claim 3, wherein the cell is an ependymal cell.
5. The method of claim 3, wherein the cell is an alveolar cell.
6. The method of claim 3, wherein the cell is a cerebellar cell.
7. The method of claim 2 wherein the cell selected from the group consisting of: alveolar cell, cerebellar cell, and ependymal cell.
8. The method of claim 7, wherein the cell is an ependymal cell.
9. The method of claim 7, wherein the cell is an alveolar cell.
10. The method of claim 7, wherein the cell is a cerebellar cell.

11. The method of any of claims 1-10, wherein the AAV inverted terminal repeats are AAV5 terminal repeats.





FIG. 1

*AAV Types 2 & 5 Vector and Helper Plasmid Combinations*



AAV2 & 5 Plasmid Combinations

FIG. 2




FIG. 3

==26-MAY-1999==

==NALIGN==

==PC/GENE==

\*\*\*\*\*  
 ALIGNMENT OF TWO NUCLEOTIDE SEQUENCES. \*  
 \*\*\*\*\*

he two sequences to be aligned are:

AV2CG.

otal number of bases: 4679.

AV5CG.

otal number of bases: 4652.

pen gap cost : 10

nit gap cost : 12

ie character to show that two aligned residues are identical is ':'

```

AV2CG - TTGGCCACTCCCTCTCTGGGCGCTCGCTCGCTEACTGA-----GGCCGGGGCGA -48
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV5CG - TGGCACTCTCCCCCTGTCTGCGCTTCGCTCGCTCGCTGGCTCGTTTGGGGGGGTGG -55
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV2CG - C-----CAAAGGTC-GCCCCACGCCCGGGCTTTGCCCCGG-GCGGCCTCA----- -90
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV5CG - CAGCTCAAAGAGCTGCCAGACGACGGCCCTCTGGCCGTGCCCCCCCCAAACGAGC -110
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV2CG - --GTGAGCGAGCGAGCGCG-CAGAGAGG-GAGTGGCCAACTCCATCACTAGGGGT -141
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV5CG - CAGCGAGCGAGCGAACGCGACAGGGGGGAGAGTGCCACACTCTCAAGCAAGGGGG -165
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV2CG - TCCTGGAGGG-GTGGAGTCGTGACG-TGAATTACGTCATAGGGTTAGGGAGGTCC -194
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV5CG - TTTTGTAAGCAGTGATGTCTAATGATGTAATGCTTATTGTACGCGATAGTTAA -220
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV2CG - TGTATTAGAGGTCACGTGA-GTGTTTTTCGACATTTTTCGACACC-----ATGT -242
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV5CG - TG-ATTAACAGTCATGTGATGTGTTTTATCCAATAGGAAGAAAGCGCGCGTATGA -274
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV2CG - GGTACAGCT-----GGGTATTTAAGCCCGAGTGAGCACGCAGGGTCTCCAT -288
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV5CG - GTTCTCGCGAGACTTCCGGGGTATAAAAGACCGAGTGAACGAGCCCCGC-CGCCAT -328
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV2CG - T-TTGAAGCGGGAG-GTTTGAACGCGCA-GCCGCCATGCCGGGGTTTACGAGAT -340
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV5CG - TCTTTGCTCTGGACTGCTAGAGGACCCCTCGCTGCCATGGCTACCTTCTATGAAGT -383
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV2CG - TGTGATTAAGGTCCCCAGCGACCTTGACGGGCATCTGCCCGGCATTTCTGACAGC -395
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV5CG - CATTGTTTCGCGTCCCATTGACGTGGAGGAACATCTGCCTGGAATTTCTGACAGC -438
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV2CG - TTTGTGAAC TGGGTGGCCGAGAAGGAATGGGAGTTGCCGCCAGATTCTGACATGG -450
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AV5CG - TTTGTGGACTGGGGTAACTGGTCAAATTTGGGAGCTGCCTCCAGAGTCAGATTTAA -493
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
  
```

FIG. 4

AV2CG - ATCTGAATCTGATGAGCAGGCACCCCTGACCGTGGCCGAGAGCTGCAGCCGA -505  
 :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::  
 AV5CG - ATTTGACTCTGGTTGAACAGCCTCAGTTGACGGTGGCTGATAGAATTGCGCCGCT -548  
 AV2CG - CTTTCTGACGGAATGGCCCGTGTGAGTAAGCCCCGGAGGCCCTTTCTTTGTG -560  
 :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::  
 AV5CG - GTTCTGTACGAGTGGAAACAAATTTTCCAAG---CAGGAGTCCAAATCTTTGTG -600  
 AV2CG - CAATTTGAGAAGGGAGAGAGCTACTTCCACATGCACGTGCTCGTGGAACCACCG -615  
 :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::  
 AV5CG - CAGTTTGAAAAGGGATCTGAATATTTTCATCTGCACACGCTTGTGGAGACCTCCG -655  
 AV2CG - GGGTGAAATCCATGGTTTGGGACGTTTCCTGAGTCAGATTCGCGAAAACTGAT -670  
 : : :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::  
 AV5CG - GCATCTCTCCATGGTCCTCGGCCGTACGTGAGTCAGATTCGCGCCAGCTGGT -710  
 AV2CG - TCAGAGAATTTACCGCGGGATCGAGCCGACTTTGCCAAACTGGTTCGCGGTCACA -725  
 :  
 AV5CG - GAAAGTGGTCTTCCAGGAATTGAACCCAGATCAACGACTGGGTGCGCCATCACC -765  
 AV2CG - AAGACCAGAAATGGCGCCGGAGGCGGGAACAAGGTGGTGATGAGTGCTACATCC -780  
 ::  
 AV5CG - AAGGTAAAGAGGGC---GGAGCC---AATAAGGTGGTGATTCTGGGTATATTC -814  
 AV2CG - CCAATTACTTGTCTCCCAAAACCAGCCTGAGCTCCAGTGGGCGTGACTAATAT -835  
 :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::  
 AV5CG - CCGCCTACCTGCTGCCGAAGGTCCAACCGAGCTTCAGTGGGCGTGGAACAACT -869  
 AV2CG - GGAACAGTATTTAAGCGCTGTGTTGAATCTCACGGAGCGTAAACGGTTGGTGGCG -890  
 :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::  
 AV5CG - GGACGAGTATAAATTGGCCGCCCTGAATCTGGAGGAGCGCAAACGGCTCGTCGCG -924  
 AV2CG - CAGCATCTGACGCACGTGTGCGAGACGAGGAGCAGAACAAGAGAATCAGAATC -945  
 :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::  
 AV5CG - CAGTTTCTGGCAGAATCCTCGCAG-CGCTCG--CAGGAGGCGGCTTCGCAGCGTG -976  
 V2CG - CCAATTCTGATGCGCCGGTGATCAGATCAAAAACCTTCAGCCAGGTACATGGAGCT -1000  
 :: : : : : :: :: :: :: : : : : : : : : : : : :  
 V5CG - AGTTCCTCGGCTGACCCGGTCATCAAAAGCAAGACTTCCAGAAATACATGGCGCT -1031  
 V2CG - GGTCCGGTGGCTCGTGGACAAGGGGATTACCTCGGAGAGCAGTGGATCCAGGAG -1055  
 ::  
 V5CG - CGTCAACTGGCTCGTGGAGCACGGCATCACTTCCGAGAAGCAGTGGATCCAGGAA -1086  
 V2CG - GACCAGGCCTCATACATCTCCTTCAATGCGGCCTCCAACCTCGCGTCCCAAATCA -1110  
 :  
 V5CG - AATCAGGAGAGCTACCTCTCCTTCAACTCCACCGGCAACTCTCGGAGCCAGATCA -1141  
 V2CG - AGGCTGCCTTGGACAATGCGGGAAGATTATGAGCCTGACTAAAACCGCCCCGA -1165  
 ::  
 V5CG - AGGCCGCGCTCGACAACGCGACCAAAATTATGAGTCTGACAAAAGCGCGGTGGA -1196  
 V2CG - CTACCTGGTGGGCCAGCAGCCCGTG-GAGGACATTTCCAGCAATCGGATTTATAA -1219  
 :: :: :: :: :: : : : : : : : : : : : : : : : :  
 V5CG - CTACCTCGTGGGG-AGCTCCGTTCCCGAGGACATTTCAAAAACAGAATCTGGCA -1250

6/34



7/34

8/34

AV2CG - CGTTCCTTTCCACAGCAGCTACGCTCACAGCCAGAGTCTGGACCGTCTCATGAAT -3507  
 ::  
 AV5CG - GGTGCCCTTCCACTCCAGCTTCGCTCCCAGTCAGAACCCTGTTCAAGCTGGCCAAC -3495  
 AV2CG - CCTCTCATCGACCAGTACCTGTATTACTT---GAGCAGAACAAACACTC----- -3553  
 :: :: : :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::  
 AV5CG - CCGCTGGTGGACCAGTACTTGTACCGCTTCGTGAGCACAAATAACACTGGCGGAG -3550  
 AV2CG - -CAAGTGAACCACCAC---GCAGTCA-AGGCTTCAGTT--TTCTCAGGCCGGAG -3601  
 : :: : ::  
 AV5CG - TCCAGTTCAACAAGAACCTGGCCGGGAGATACGCCAACACCTACAAAACTGGTT -3605  
 AV2CG - CGAGTGACATTCCGGACCAGTCTAGGAACTGGCTTCCTGGACCCCTGTTACCGCCA -3656  
 :  
 AV5CG - CCCGGGGCCCATGGGCCGAACCCAGGG-CTGGAA-CCTGGGCTCCGGGTCAACC -3658  
 AV2CG - GCAGCGAGTATCAAAGACATCTGCGGATAACAACAACAGTGAATACTCGTGGACT -3711  
 :: :: :: :: ::  
 AV5CG - GC-GCCAGTGTGAGCGCCTTC-GCCACGACCAATAGGA-TGGAG-CTCGAGGGCG -3709  
 AV2CG - GGAGCTACCAAGTACCACCTCAATGGCAGAGACTCTCTGGTGAATCCGGGCCCGG -3766  
 :: :: :: :: ::  
 AV5CG - CGAGTTACCAGGTGCCCCCGCA---GCGA-ACGGCATGACCAACAACCTCCAGG -3760  
 AV2CG - CCATGGCAAGCCACAAGGACGATGAAGAAAGTTTTCCTCAGAGCGGGTTCT -3821  
 :: :: :: :: ::  
 AV5CG - GCA--GCAA--CACCTATGCCCTGGAGAACACTATGATCTTCAA--CAG----C- -3804  
 AV2CG - CATCTTTGGGAAGCAAGGCTCAGAGAAAACAAATGTGGACATTGAAAAGGTCATG -3876  
 ::  
 AV5CG - CAGCCG-GCGAACCCGGGCACCACCGCCACGTACCTCGAGGGCAACATGCTCATC -3858  
 AV2CG - ATTACAGACGAAGAGGAAATCAGGACAACCAATCCCGTGGC-TACGGAGCAGTAT -3930  
 : :: : ::  
 AV5CG - AC--CAG-CGAGAGCGAGACGCAGCCGGTGAACCGCGTGGCGTACAACGTCCGCG -3910  
 AV2CG - GGTTCGTATCTACCAACCTCCAGAGAGGCAACAGACAAGCAGCTACCGCAGATG -3985  
 :  
 AV5CG - GGCAGA-TGGCCACCAACAACCAGAGCTCCACCACTGCCCCCGCGACCGGCACGT -3964  
 AV2CG - TCAACACACAAGGCGTTCTTCAGGCATGGTCTGGCAGGACAGAGATGTGTACCT -4040  
 :: ::  
 AV5CG - ACAACCTCCAGGAAATCGTGCCCGGCAGCGTGTGGATGGAGAGGGACGTGTACCT -4019  
 AV2CG - TCAGGGGCCCATCTGGGCAAAGATTCCACACACGGACGGACATTTTCACCCCTCT -4095  
 : : : : :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::  
 AV5CG - CCAAGGACCCATCTGGGCCAAGATCCCAGAGACGGGGGCGCACTTTTCACCCCTCT -4074  
 AV2CG - CGCCTCATGGGTGGATTGGACTTAAACACCCCTCCTCCACAGATTCTCATCAAGA -4150  
 :: : :: :: :: :: :: :: :: :: :: :: :: : : : : : : : : : :  
 AV5CG - CCGGCCATGGGCGGATTCGGACTCAAAACCCACCGCCCATGATGCTCATCAAGA -4129  
 AV2CG - ACACCCCGGTACCTGCGAATCCTTCGACCACCTTCAGTG-CGGCAAAGTTTGCTT -4204  
 :: ::  
 AV5CG - ACACGCCTGTGCCCGGAAATA--TC-ACCAGCTTCTCGGACGTGCCCGTCAGCAG -4181

```

ntity : 3013 (64.77%)
ber of gaps inserted in AAV2CG: 43
ber of gaps inserted in AAV5CG: 63

```

26-MAY-1999=====NATIGN=====PC/GENE=====

22-JAN-1997 PALIGN PC/GENE

\*\*\*\*\*  
 \* ALIGNMENT OF TWO PROTEIN SEQUENCES. \*  
 \*\*\*\*\*

The two sequences to be aligned are:

TRP 6/AC 6

AAV2VP1.  
 DE VP1  
 OS AAV2  
 Total number of residues: 735.

AAV5VP1.  
 DE AAV5VP1  
 OS AAV5VP1  
 Total number of residues: 724. 2/01-4272

Comparison matrix : Structure-genetic matrix.  
 Open gap cost : 8  
 Unit gap cost : 5

The character to show that two aligned residues are identical is ':'  
 The character to show that two aligned residues are similar is '.'  
 Amino acids said to be 'similar' are: A,S,T; D,E; N,Q; R,K; I,L,M,V; F,Y,W

```

AAV2VP1 - MAADGYLPDWLEDTLSEGIQWWKLPKPPPKPAERHKDDSRGLVLPGYKYLGP -55
          ::::::::::: . : : : : : : : : : : : : : : : : : : : : : : : :
AAV5VP1 - MSFVDHPPDWLEE-VGEGLREFLGLEAGPKPKPNQOHQDQARGLVLPGYNYLGP -54
          ::::::::::: . : : : : : : : : : : : : : : : : : : : : : : : :

AAV2VP1 - ENGLDKGEPVNEADAAALEHDKAYDRQLDSDGNPYLKYNHADAEFQERLKEDTSF -110
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :
AAV5VP1 - GNGLDRGEPVNRADDEVAREHDISYNEQLEAGDNPYLKYNHADAEFQEKLADDTSF -109
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :

AAV2VP1 - GGNLGRAVFQAKKRVLEPLGLVEEPVKTPAGKKRPVEHSPVEPDSSSGTGKAGQQ -165
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :
AAV5VP1 - GGNLGKAVFQAKKRVLEPFGLVEEGAKTAPTGKRIDDHFPKR--KKARTEEDSKP -162
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :

AAV2VP1 - PARKRLNFGQTGDADSVDPDQPLGQPPAAPSGLGTTMTATGSGAPMADNNEGADG -220
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :
AAV5VP1 - STS-----SDAEAGPSGSQQLQIPAQPASSLGADTMSAGGGGPLGDNNOGADG -210
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :

AAV2VP1 - VGNSSGNWHCDSTWMGDRVITTTSTRTWALPTYNNHLYKQISSQSG-ASNDNHYFG -274
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :
AAV5VP1 - VGNASGDWHCDSTWMGDRVVTKSTRTWVLPSTNNHLYKQREIKSGSVDGSNANAYFG -265
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :

AAV2VP1 - YSTPWGYFDNRFHCHFSFRDWQRLINNNWGFPRPKRLNFKLENIQVKEVTQNDGT -329
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :
AAV5VP1 - YSTPWGYFDNRFHSHWSFRDWQRLINNYWGFPRPSLRVKIFNIQVKEVTVQDST -320
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :

AAV2VP1 - TTIANNLTSTVQVFTDSEYQLPYVLGSAHQGCLPPFFADVFMVPOYGYLTLNNGS -384
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :
AAV5VP1 - TTIANNLTSTVQVFTDDDYQLPYVVGNGTEGCLPAFFPPQVFTLPQYGYATLNDRN -375
          ::::::::::: . : : : . : : : : : : : : : : : : : : : : : : : :
  
```

```

AAV2VP1 - Q--AVGRSSFYCLEYFPSQMLRTGNNFTFSYTFEDVPFESSYAHSQSLDRIMNPL -437
          :::::::::::::::::::: :: :::::::::::::: :: :: ::
AAV5VP1 - TENPTERSSFFCLEYFPSKMLRTGNNFEFTYNFEEVPFHSSFAPSQNLFKLANPL -430
AAV2VP1 - IDQYLYYLSRTNTPSGTTTQSRLOFSQAGASDIRDQSRNWLPGPCYRQQRVSKTS -492
          ::::: :: :: ::::: ::::: ::::: ::::: :::::
AAV5VP1 - VDQYLYRFVSTNNTGG-----VQFNKNLAGRYANTYKNWFPGPMGRTQGWNLGS -479
AAV2VP1 - ADNNNSEYSWGATKYHLNGRDSLVPNGPAMASHKDDEEKFFPQSGVLIFGKQGS -547
          : : . . . . . : : : : : : : : : : : :
AAV5VP1 - GVNRAVSASFATTNRMELEGASYQVPPQPNGMTNNLQGSNTYALENTMIFNSQPA -534
AAV2VP1 - EKINVDI---EKVMITDEEEIIRTTNPVATEQYGSVSTNLQRGNRQAATADVNTQG -599
          . . . : : : : : : : : : : : : : :
AAV5VP1 - NPGTTATYLEGNMLITSESETQPVNRVAYNVGGQMATNNQSSTTAPATGTYNLQE -589
AAV2VP1 - VLPGMVWQDRDVYLQGPWAKIPETDGHFHPSPLMGGFGLKHPPPPQILIKNTFVP -654
          . . : : . . . . . : : : : : : : : : : : :
AAV5VP1 - IVPGSVWMERDVYLQGPWAKIPETGAHFHPSPAMGGFGLKHPPPPMMLIKNTFVP -644
AAV2VP1 - ANPSTTFSAAKFASFITQYSTGQVSVEIEWELQKENS KRWNPEIQYTSNYNKSVN -709
          : : . . . . . : : : : : : : : : : : :
AAV5VP1 - GNI-TSFSQVPSVSSFITQYSTGQVTVEMEWELKKENS KRWNPEIQYTNNYNQPF -698
AAV2VP1 - VDFTVDTNGVYSEPRPIGTRYLTRNL -735
          : . . : : : : : : : : :
AAV5VP1 - VDFAPDSTGEYRTRPIGTRYLTRPL -724

```

Identity : 421 (58.15%)

Similarity: 63 (8.70%)

Number of gaps inserted in AAV2VP1: 3

Number of gaps inserted in AAV5VP1: 5

==22-JAN-1997==PALIGN==PC/GENE==

## FIG. 6

31-DEC-1996 PC/GENE

\*\*\*\*\*  
 \* ALIGNMENT OF TWO PROTEIN SEQUENCES. \*  
 \*\*\*\*\*

The two sequences to be aligned are:

REP78.

DE REP78

OS AAV

Total number of residues: 621.

AAV5REP.

DE REP

OS AAV5

Total number of residues: 610.

Comparison matrix : Structure-genetic matrix.

Open gap cost : 8

Unit gap cost : 5

The character to show that two aligned residues are identical is ':'

The character to show that two aligned residues are similar is '.'

Amino acids said to be 'similar' are: A,S,T; D,E; N,Q; R,K; I,L,M,V; F,Y,W

```

REP78      - MPGFYIEIVIKVPSDLGDGHLPGISDSFVNWVAEKELPPDSMDLNLIEQAPLTV -55
              : ::::: : : : : : : : : : : : : : : : : : : : : : : : : : :
AAV5REP    - MATFYEVIVRVFPDVEEHLPGISDSFVDWVTGQIWELPPESDLNLTLVEQPQLTV -55

REP78      - AEKLORDFLTEWRRVSKAPEALFFVQFEKGSYFMEVLVETTGVKSMVLGRFLS -110
              : : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AAV5REP    - ADRIIRRVFLYEWNKFSKQ-ESKFFVQFEKGSYFHELTIVETSGISSMVLGRYVS -109

REP78      - QIREKLIQRIYRGIEPTLPNWFAVTKTRNGAGGKNKVVDDECYIPNYLLPKTQPEL -165
              : : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AAV5REP    - QIRAOVLVKVVFQGLEPQINDWVAITKVKKG--GANKVVDSCYIPAYILLEPKVQPEL -162

REP78      - QWAWTNMEQYLSACLNLTERRKRLVAQHLTHVSQTQEONKENQNPNSDAPVIRSKT -220
              : : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AAV5REP    - QWAWTNLDEYKLAALNLEERKRLVAQFLA-ESSQRSQEAASQREFSADPVIKSKT -216

REP78      - SARYMELVGWLVDKGITSEKQWIQEDQASYISFNAASNSRSQIKAALDNACKIMS -275
              : : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AAV5REP    - SQKYMALVNWLVEHGITSEKQWIQENQESYLSFNSTGNSRSQIKAALDNATKIMS -271

REP78      - LTKTAPDYLVGQQPVEDISSNRIYKILELNGYDPOQYASVFLGWATKKFGRNTI -330
              : : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AAV5REP    - LTKSAVDYLVGSSVPEDLSKNRIWQIFEMNGYDPAYAGSILYGWCQORSFNKRNTV -326

REP78      - WLFGPATTGKTNTIAEAIHTVPPFYGCNVNWTNENFPFND CVDKMWIWWEEGKMTAK -385
              : : : : : : : : : : : : : : : : : : : : : : : : : : : : :
AAV5REP    - WLYGPATTGKTNTIAEAIHTVPPFYGCNVNWTNENFPFND CVDKMLIWWEEGKMINK -381
  
```

```

REP78      - VVESAKAILGGSKVRVDQKCKSSAQIDPTPVIVTSNTNMCVIDGNSTTFEEQQP -440
             :::::::::::::::::::::: :: :::::::::::::: ::::::::::::::
AAV5REP    - VVESAKAILGGSKVRVDQKCKSSVQIDSTPVIVTSNTNMCVVVDGNSTTFEEQQP -436
             :::::::::::::::::::::: :: :::::::::::::: ::::::::::::::
REP78      - LQDRMFKFELTRRLDHDGKVTKEVVDKDFRWAKDHVVEVEHEFYVKKGCAKKRP -495
             : :::::::::::::: :::::::::::::: :: ::::::::::::::
AAV5REP    - LEDRMFKFELTKRLPPDFGKITKQEVKDEFFAWAKVNQVPVTHEFKVPRELAKTK- -490
             :::::::::::::: :::::::::::::: ::::::::::::::
REP78      - APSDADISEPKRVRESVAQPSTSDAEASINYADRYQNKCSRHVGMNLMLEFCRQC -550
             : :::::::::::::: ::::::::::::::
AAV5REP    - GAEKSLKRPLGDVTNTXYKSLEKRRLSFVPETPRSSDVTVDPAFLRPLNWNRSRY -545
             : :::::::::::::: ::::::::::::::
REP78      - ERMNQNSNICFTEGQKDCLECFPVSESQPVSVVKKAYQKLCYIEHIMGKVPDACT -605
             : :::::::::::::: ::::::::::::::
AAV5REP    - DC-KCDYHAQFDNISNKCDECEYLNRGKNGCICHNVTH-CQICHGIPPWEKENLS -598
             : ::::::::::::::
REP78      - ACDLVNVDLDDCIFEQ -621
             : :: :: ::
AAV5REP    - DF----GDFDDANKEQ -610

```

Identity : 355 ( 58.2%)

Similarity: 56 ( 9.2%)

Number of gaps inserted in REP78: 0

Number of gaps inserted in AAV5REP: 7

==31-DEC-1996==

PC/GENE



Apical transduction of human airway epithelia with rAAV2 and rAAV5




FIG. 7





FIG. 8

rAAV5 Primary Rat Brain Explant



**FIG. 9**

HUVEC

rAAV2

rAAV5



**FIG. 10**




Fig. 11.




Fig. 12




Fig. 13




Fig. 14






Fig. 15




Fig. 16A





Fig. 16B




Fig. 17




Fig 18




Fig. 19A




Fig. 19 B




Fig 20 A + B






Fig 20C




Fig. 21




Fig. 22




Fig 23

## SEQUENCE LISTING

<110> The Government of the United States of America, as  
represented by the Secretary, Department of Health and Human  
Services

Chiorini, John  
Kotin, Robert M.  
Davidson, Elizabeth  
Zabner, Joseph

<120> AAV5 VECTOR FOR TRANSDUCING BRAIN CELLS AND LUNG CELLS

<130> 14014.0323P2

<140> 09/533,427

<141> 2000-03-22

<160> 23

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 4652

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/Note =  
synthetic construct

<400> 1

tggaactctc	ccccctgtcg	cgttcgctcg	ctcgctggct	cgtttggggg	ggtggcagct	60
caaagagctg	ccagacgacg	gccctctggc	cgtcgcccc	ccaaacgagc	cagcgagcga	120
gcgaacgcga	caggggggag	agtgccacac	tctcaagcaa	gggggttttg	taagcagtga	180
tgtcataatg	atgtaatgct	tattgtcagc	cgatagttaa	tgattaacag	tcatgtgatg	240
tgttttatcc	aataggaaga	aagcgcgcgt	atgagttctc	gcgagacttc	cgggggtataa	300
aagaccgagt	gaacgagccc	gccgccattc	tttgctctgg	actgctagag	gaccctcgct	360
gccatggcta	ccttctatga	agtcattgtt	cgcgtcccat	ttgacgtgga	ggaacatctg	420
cctggaattt	ctgacagctt	tgtggactgg	gtaactggtc	aaatttggga	gctgcctcca	480
gagtcagatt	taaatgtgac	tctgggtgaa	cagcctcagt	tgacggtggc	tgatagaatt	540
cgccgcgtgt	tcctgtacga	gtggaacaaa	ttttccaagc	aggagtccaa	attctttgtg	600
cagtttgaaa	agggatctga	atattttcat	ctgcacacgc	ttgtggagac	ctccggcatc	660
tcttccatgg	tcctcggcgc	ctacgtgagt	cagattcgcg	cccagctggg	gaaagtggtc	720
ttccaggga	ttgaacccca	gatcaacgac	tgggtcgcca	tcaccaaggt	aaagaagggc	780
ggagccaata	aggtgggtga	ttctgggtat	attccgcct	acctgctgcc	gaagggtcaa	840
ccggagcttc	agtgggcgtg	gacaaacctg	gacgagtata	aattggccgc	cctgaatctg	900
gaggagcgca	aacggctcgt	cgcgcagttt	ctggcagaat	cctcgacgag	ctcgagggag	960
gcggcttcgc	agcgtgagtt	ctcggtcgac	ccggtcatca	aaagcaagac	ttcccagaaa	1020
tacatggcgc	tcgtcaactg	gctcgtggag	cacggcatca	cttccgagaa	gcagtggatc	1080
caggaataatc	aggagagcta	cctctccttc	aactccaccg	gcaactctcg	gagccagatc	1140
aaggccgcgc	tcgacaacgc	gaccaaaatt	atgagttctg	caaaaagcgc	ggtggactac	1200
ctcgtgggga	gtcccggttc	cgaggacatt	tcaaaaaaca	gaatctggca	aatttttgag	1260
atgaatggct	acgacccggc	ctacgcggga	tccatcctct	acggctggtg	tcagcgctcc	1320
ttcaacaaga	ggaacaccgt	ctggctctac	ggaccgcgca	cgaccggcaa	gaccaacatc	1380
gcggaggcca	tcgcccacac	tgtgcccttt	tacggctgcg	tgaactggac	caatgaaaac	1440
tttcccttta	atgactgtgt	ggacaaaatg	ctcatttggg	gggaggaggg	aaagatgacc	1500
aacaaggtgg	ttgaatccgc	caaggccatc	ctggggggct	caaagggtgcg	ggtcgatcag	1560
aatgtaaat	cctctgttca	aattgattct	accctgttca	ttgtaacttc	caatacaaac	1620

atgtgtgtgg	tgggtggatgg	gaattccacg	acctttgaac	accagcagcc	gctggaggac	1680
cgcagtgtca	aattttgaact	gactaagcgg	ctcccgccag	attttggcaa	gattactaag	1740
caggaagtca	aggacttttt	tgcttgggca	aaggtcaatc	aggtgccggt	gactcacgag	1800
tttaaagttc	ccaggggaatt	ggcgggaact	aaaggggcgg	agaaatctct	aaaacgcca	1860
ctgggtgacg	tcaccaatac	tagctataaa	agtctggaga	agcggggccag	gctctcattt	1920
gttccccgaga	cgcctcgcag	ttcagacgtg	actgttgatc	ccgctcctct	gcgaccgctc	1980
aattggaatt	caaggtatga	ttgcaaattg	gactatcatg	ctcaatttga	caacatttct	2040
aacaaatgtg	atgaatgtga	atattttgaat	cggggcaaaa	atggatgtat	ctgtcacaat	2100
gtaactcact	gtcaaatttg	tcattgggatt	ccccctggg	aaaaggaaaa	cttgtcagat	2160
tttggggatt	ttgacgatgc	caataaagaa	cagtaaataa	agcgagtagt	catgtctttt	2220
gttgatcacc	ctccagattg	gttgggaagaa	gttgggtgaag	gtcttcgcca	gtttttgggc	2280
cttgaagcgg	gcccaccgaa	acaaaaaccc	aatcagcagc	atcaagatca	agcccggtgt	2340
cttgtgctgc	ctgggtataa	ctatctcgga	cccggaacg	gtctcgatcg	aggagagcct	2400
gtcaacaggg	cagacgaggt	cgcgcgagag	cacgacatct	cgtacaacga	gcagcttgag	2460
gcggggagaca	acccctacct	caagtacaac	cacgcggacg	ccgagtttca	ggagaagctc	2520
gccgacgaca	cactcttcgg	gggaaacctc	ggaaaggcag	tctttcaggc	caagaaaagg	2580
gttctcgaac	cttttggcct	ggttgaagag	ggtgctaaga	cggccccctac	cggaaagcgg	2640
atagacgacc	actttccaaa	aagaaagaag	gctcggaccg	aagaggactc	caagccttcc	2700
acctcgtcag	acgccgaagc	tggacccagc	ggatcccagc	agctgcaaat	cccagcccaa	2760
ccagcctcaa	gttttgggagc	tgatacaatg	tctgcgggag	gtggcgggcc	attggggcag	2820
aataaccaag	gtgccgatgg	agtgggcaat	gcctcgggag	attggcattg	cgattccacg	2880
tggatggggg	acagagtcgt	caccaagtcc	acccgaacct	gggtgctgcc	cagctacaac	2940
aaccaccagt	accgagagat	caaaagcggc	tccgtcgacg	gaagcaacgc	caacgcctac	3000
tttgatatac	gcacccctcg	ggggtaactt	gactttaacc	gcttccacag	ccactggagc	3060
ccccgagact	ggcaaagact	catcaacaac	tactggggct	tcagaccccg	gtccctcaga	3120
gtcaaaatct	tcaacattca	agtcaaagag	gtcacgggtg	aggactccac	caccaccatc	3180
gccaacaacc	tcacctccac	cgtccaagtg	tttacggacg	acgactacca	gctgccttac	3240
gtcgtcggca	acgggaccga	gggatgcctg	ccggccttcc	ctccgcaggt	ctttacgctg	3300
ccgcagtacg	gttacgcgac	gctgaaccgc	gacaacacag	aaaatcccac	cgagaggagc	3360
agcttcttct	gcctagagta	ctttcccagc	aagatgctga	gaacgggcaa	caactttgag	3420
tttacctaca	actttgagga	ggtgcccttc	cactccagct	tcgctcccag	tcagaacctg	3480
ttcaagctgg	ccaaccgct	ggtggaccag	tacttgtacc	gcttcgtgag	cacaaataac	3540
actggcgagg	tcagttcaa	caagaacctg	ccggggagat	acgccaacac	ctacaaaaac	3600
tgggttcccg	ggcccatggg	ccgaacccag	ggctggaacc	tgggctccgg	ggtcaaccgc	3660
gccagtgtca	gcgccttcgc	cacgaccaat	aggatggagc	tcgaggggcg	gagttaccag	3720
gtgcccccg	agccgaacgg	catgaccaac	aacctccagg	gcagcaaac	ctatgccctg	3780
gagaacacta	tgatcttcaa	cagccagccg	gcgaacccgg	gcaccaccgc	cagtaacctc	3840
gagggcaaca	tgctcatcac	cagcgagagc	gagacgcagc	cgttgaaccg	cgtggcgta	3900
aacgtcggcg	ggcagatggc	caccaacaac	cagagctcca	ccactgcccc	cgcgaccggc	3960
acgtacaacc	tccaggaaat	cgtgcccggc	agcgtgtgga	tggagaggga	cgtgtacctc	4020
caaggaccca	tctggggcaa	gatcccagag	acggggggcg	actttcaccc	ctctccggcc	4080
atggggggat	tcggactcaa	acacccaccg	cccatgatgc	tcatacaaga	cacgcctgtg	4140
cccggaaata	tcaccagctt	ctcggacgtg	cccgtcagca	gcttcatcac	ccagtacagc	4200
accgggcagg	tcaccgtgga	gatggagtgg	gagctcaaga	aggaaaaact	caagaggtgg	4260
aaccagagaa	tccagtacac	aaacaactac	aacgaccccc	agttttgtgga	ctttgccccg	4320
gacagcaccg	gggaatacag	aaccaccaga	cctatcgga	cccgatacct	taccgacccc	4380
ctttaaccca	ttcatgtcgc	ataccctcaa	taaaccgtgt	attcgtgtca	gtaaaaatac	4440
gcctcttctg	gtcatttcaat	gaataacagc	ttacaacatc	tacaaaaact	ccttgcttga	4500
gagtgtggca	ctctcccccc	tgtcgcgttc	gctcgcctcg	tggctcgttt	gggggggtgg	4560
cagctcaaag	agctgccaga	cgacggccct	ctggccgctg	cccccccaaa	cgagccagcg	4620
agcgagcgaa	cgcgacaggg	gggagagtgc	ca			4652

&lt;210&gt; 2

&lt;211&gt; 390

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
synthetic construct

&lt;400&gt; 2

Met Ala Leu Val Asn Trp Leu Val Glu His Gly Ile Thr Ser Glu Lys

3

```

1           5           10           15
Gln Trp Ile Gln Glu Asn Gln Glu Ser Tyr Leu Ser Phe Asn Ser Thr
20           25           30
Gly Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Thr Lys
35           40           45
Ile Met Ser Leu Thr Lys Ser Ala Val Asp Tyr Leu Val Gly Ser Ser
50           55           60
Val Pro Glu Asp Ile Ser Lys Asn Arg Ile Trp Gln Ile Phe Glu Met
65           70           75
Asn Gly Tyr Asp Pro Ala Tyr Ala Gly Ser Ile Leu Tyr Gly Trp Cys
85           90           95
Gln Arg Ser Phe Asn Lys Arg Asn Thr Val Trp Leu Tyr Gly Pro Ala
100          105          110
Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Thr Val Pro
115          120          125
Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp
130          135          140
Cys Val Asp Lys Met Leu Ile Trp Trp Glu Glu Gly Lys Met Thr Asn
145          150          155
Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg
165          170          175
Val Asp Gln Lys Cys Lys Ser Ser Val Gln Ile Asp Ser Thr Pro Val
180          185          190
Ile Val Thr Ser Asn Thr Asn Met Cys Val Val Val Asp Gly Asn Ser
195          200          205
Thr Thr Phe Glu His Gln Gln Pro Leu Glu Asp Arg Met Phe Lys Phe
210          215          220
Glu Leu Thr Lys Arg Leu Pro Pro Asp Phe Gly Lys Ile Thr Lys Gln
225          230          235
Glu Val Lys Asp Phe Phe Ala Trp Ala Lys Val Asn Gln Val Pro Val
245          250          255
Thr His Glu Phe Lys Val Pro Arg Glu Leu Ala Gly Thr Lys Gly Ala
260          265          270
Glu Lys Ser Leu Lys Arg Pro Leu Gly Asp Val Thr Asn Thr Ser Tyr
275          280          285
Lys Ser Leu Glu Lys Arg Ala Arg Leu Ser Phe Val Pro Glu Thr Pro
290          295          300
Arg Ser Ser Asp Val Thr Val Asp Pro Ala Pro Leu Arg Pro Leu Asn
305          310          315
Trp Asn Ser Arg Tyr Asp Cys Lys Cys Asp Tyr His Ala Gln Phe Asp
325          330          335
Asn Ile Ser Asn Lys Cys Asp Glu Cys Glu Tyr Leu Asn Arg Gly Lys
340          345          350
Asn Gly Cys Ile Cys His Asn Val Thr His Cys Gln Ile Cys His Gly
355          360          365
Ile Pro Pro Trp Glu Lys Glu Asn Leu Ser Asp Phe Gly Asp Phe Asp
370          375          380
Asp Ala Asn Lys Glu Gln
385          390

```

```

<210> 3
<211> 610
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence:/Note =
        synthetic construct

```

```

<400> 3
Met Ala Thr Phe Tyr Glu Val Ile Val Arg Val Pro Phe Asp Val Glu
1           5           10           15
Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Asp Trp Val Thr Gly

```

				20					25				30			
Gln	Ile	Trp	Glu	Leu	Pro	Pro	Glu	Ser	Asp	Leu	Asn	Leu	Thr	Leu	Val	
		35					40					45				
Glu	Gln	Pro	Gln	Leu	Thr	Val	Ala	Asp	Arg	Ile	Arg	Arg	Val	Phe	Leu	
	50					55					60					
Tyr	Glu	Trp	Asn	Lys	Phe	Ser	Lys	Gln	Glu	Ser	Lys	Phe	Phe	Val	Gln	
65				70						75					80	
Phe	Glu	Lys	Gly	Ser	Glu	Tyr	Phe	His	Leu	His	Thr	Leu	Val	Glu	Thr	
			85						90					95		
Ser	Gly	Ile	Ser	Ser	Met	Val	Leu	Gly	Arg	Tyr	Val	Ser	Gln	Ile	Arg	
			100					105					110			
Ala	Gln	Leu	Val	Lys	Val	Val	Phe	Gln	Gly	Ile	Glu	Pro	Gln	Ile	Asn	
		115					120					125				
Asp	Trp	Val	Ala	Ile	Thr	Lys	Val	Lys	Lys	Gly	Gly	Ala	Asn	Lys	Val	
	130					135					140					
Val	Asp	Ser	Gly	Tyr	Ile	Pro	Ala	Tyr	Leu	Leu	Pro	Lys	Val	Gln	Pro	
145				150						155					160	
Glu	Leu	Gln	Trp	Ala	Trp	Thr	Asn	Leu	Asp	Glu	Tyr	Lys	Leu	Ala	Ala	
			165						170					175		
Leu	Asn	Leu	Glu	Glu	Arg	Lys	Arg	Leu	Val	Ala	Gln	Phe	Leu	Ala	Glu	
			180					185					190			
Ser	Ser	Gln	Arg	Ser	Gln	Glu	Ala	Ser	Gln	Arg	Glu	Phe	Ser	Ala		
		195					200				205					
Asp	Pro	Val	Ile	Lys	Ser	Lys	Thr	Ser	Gln	Lys	Tyr	Met	Ala	Leu	Val	
	210					215					220					
Asn	Trp	Leu	Val	Glu	His	Gly	Ile	Thr	Ser	Glu	Lys	Gln	Trp	Ile	Gln	
225				230						235					240	
Glu	Asn	Gln	Glu	Ser	Tyr	Leu	Ser	Phe	Asn	Ser	Thr	Gly	Asn	Ser	Arg	
			245						250					255		
Ser	Gln	Ile	Lys	Ala	Ala	Leu	Asp	Asn	Ala	Thr	Lys	Ile	Met	Ser	Leu	
			260					265					270			
Thr	Lys	Ser	Ala	Val	Asp	Tyr	Leu	Val	Gly	Ser	Ser	Val	Pro	Glu	Asp	
		275					280					285				
Ile	Ser	Lys	Asn	Arg	Ile	Trp	Gln	Ile	Phe	Glu	Met	Asn	Gly	Tyr	Asp	
	290					295					300					
Pro	Ala	Tyr	Ala	Gly	Ser	Ile	Leu	Tyr	Gly	Trp	Cys	Gln	Arg	Ser	Phe	
305				310						315					320	
Asn	Lys	Arg	Asn	Thr	Val	Trp	Leu	Tyr	Gly	Pro	Ala	Thr	Thr	Gly	Lys	
			325						330					335		
Thr	Asn	Ile	Ala	Glu	Ala	Ile	Ala	His	Thr	Val	Pro	Phe	Tyr	Gly	Cys	
			340					345					350			
Val	Asn	Trp	Thr	Asn	Glu	Asn	Phe	Pro	Phe	Asn	Asp	Cys	Val	Asp	Lys	
		355					360									



5

Lys Arg Ala Arg Leu Ser Phe Val Pro Glu Thr Pro Arg Ser Ser Asp  
           515                          520                          525  
 Val Thr Val Asp Pro Ala Pro Leu Arg Pro Leu Asn Trp Asn Ser Arg  
           530                          535                          540  
 Tyr Asp Cys Lys Cys Asp Tyr His Ala Gln Phe Asp Asn Ile Ser Asn  
   545                          550                          555                          560  
 Lys Cys Asp Glu Cys Glu Tyr Leu Asn Arg Gly Lys Asn Gly Cys Ile  
                           565                          570                          575  
 Cys His Asn Val Thr His Cys Gln Ile Cys His Gly Ile Pro Pro Trp  
                           580                          585                          590  
 Glu Lys Glu Asn Leu Ser Asp Phe Gly Asp Phe Asp Asp Ala Asn Lys  
                           595                          600                          605  
 Glu Gln  
   610

<210> 4  
 <211> 724  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
           synthetic construct

<400> 4  
 Met Ser Phe Val Asp His Pro Pro Asp Trp Leu Glu Glu Val Gly Glu  
   1                          5                          10                          15  
 Gly Leu Arg Glu Phe Leu Gly Leu Glu Ala Gly Pro Pro Lys Pro Lys  
                           20                          25                          30  
 Pro Asn Gln Gln His Gln Asp Gln Ala Arg Gly Leu Val Leu Pro Gly  
                           35                          40                          45  
 Tyr Asn Tyr Leu Gly Pro Gly Asn Gly Leu Asp Arg Gly Glu Pro Val  
   50                          55                          60  
 Asn Arg Ala Asp Glu Val Ala Arg Glu His Asp Ile Ser Tyr Asn Glu  
   65                          70                          75                          80  
 Gln Leu Glu Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala Asp  
                           85                          90                          95  
 Ala Glu Phe Gln Glu Lys Leu Ala Asp Asp Thr Ser Phe Gly Gly Asn  
                           100                          105                          110  
 Leu Gly Lys Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro Phe  
                           115                          120                          125  
 Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Thr Gly Lys Arg Ile  
   130                          135                          140  
 Asp Asp His Phe Pro Lys Arg Lys Lys Ala Arg Thr Glu Glu Asp Ser  
   145                          150                          155                          160  
 Lys Pro Ser Thr Ser Ser Asp Ala Glu Ala Gly Pro Ser Gly Ser Gln  
                           165                          170                          175  
  
 Gln Leu Gln Ile Pro Ala Gln Pro Ala Ser Ser Leu Gly Ala Asp Thr  
                           180                          185                          190  
 Met Ser Ala Gly Gly Gly Gly Pro Leu Gly Asp Asn Asn Gln Gly Ala  
                           195                          200                          205  
 Asp Gly Val Gly Asn Ala Ser Gly Asp Trp His Cys Asp Ser Thr Trp  
   210                          215                          220  
 Met Gly Asp Arg Val Val Thr Lys Ser Thr Arg Thr Trp Val Leu Pro  
   225                          230                          235                          240  
 Ser Tyr Asn Asn His Gln Tyr Arg Glu Ile Lys Ser Gly Ser Val Asp  
                           245                          250                          255  
 Gly Ser Asn Ala Asn Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr  
                           260                          265                          270  
 Phe Asp Phe Asn Arg Phe His Ser His Trp Ser Pro Arg Asp Trp Gln  
                           275                          280                          285  
 Arg Leu Ile Asn Asn Tyr Trp Gly Phe Arg Pro Arg Ser Leu Arg Val

290	295	300
Lys Ile Phe Asn Ile Gln Val	Lys Glu Val Thr Val Gln Asp Ser Thr	
305	310	315
Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp		320
	325	330
Asp Asp Tyr Gln Leu Pro Tyr Val Val Gly Asn Gly Thr Glu Gly Cys		335
	340	345
Leu Pro Ala Phe Pro Pro Gln Val Phe Thr Leu Pro Gln Tyr Gly Tyr		350
	355	360
Ala Thr Leu Asn Arg Asp Asn Thr Glu Asn Pro Thr Glu Arg Ser Ser		365
	370	375
Phe Phe Cys Leu Glu Tyr Phe Pro Ser Lys Met Leu Arg Thr Gly Asn		380
385	390	395
Asn Phe Glu Phe Thr Tyr Asn Phe Glu Glu Val Pro Phe His Ser Ser		400
	405	410
Phe Ala Pro Ser Gln Asn Leu Phe Lys Leu Ala Asn Pro Leu Val Asp		415
	420	425
Gln Tyr Leu Tyr Arg Phe Val Ser Thr Asn Asn Thr Gly Gly Val Gln		430
	435	440
Phe Asn Lys Asn Leu Ala Gly Arg Tyr Ala Asn Thr Tyr Lys Asn Trp		445
	450	455
Phe Pro Gly Pro Met Gly Arg Thr Gln Gly Trp Asn Leu Gly Ser Gly		460
465	470	475
Val Asn Arg Ala Ser Val Ser Ala Phe Ala Thr Thr Asn Arg Met Glu		480
	485	490
Leu Glu Gly Ala Ser Tyr Gln Val Pro Pro Gln Pro Asn Gly Met Thr		495
	500	505
Asn Asn Leu Gln Gly Ser Asn Thr Tyr Ala Leu Glu Asn Thr Met Ile		510
	515	520
Phe Asn Ser Gln Pro Ala Asn Pro Gly Thr Thr Ala Thr Tyr Leu Glu		525
	530	535
Gly Asn Met Leu Ile Thr Ser Glu Ser Glu Thr Gln Pro Val Asn Arg		540
545	550	555
Val Ala Tyr Asn Val Gly Gly Gln Met Ala Thr Asn Asn Gln Ser Ser		560
	565	570
Thr Thr Ala Pro Ala Thr Gly Thr Tyr Asn Leu Gln Glu Ile Val Pro		575
	580	585
Gly Ser Val Trp Met Glu Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp		590
	595	600
	605	
Ala Lys Ile Pro Glu Thr Gly Ala His Phe His Pro Ser Pro Ala Met		
610	615	620
Gly Gly Phe Gly Leu Lys His Pro Pro Pro Met Met Leu Ile Lys Asn		
625	630	635
Thr Pro Val Pro Gly Asn Ile Thr Ser Phe Ser Asp Val Pro Val Ser		
	645	650
Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Thr Val Glu Met Glu		
	660	665
Trp Glu Leu Lys Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln		
	675	680
Tyr Thr Asn Asn Tyr Asn Asp Pro Gln Phe Val Asp Phe Ala Pro Asp		
	690	695
Ser Thr Gly Glu Tyr Arg Thr Thr Arg Pro Ile Gly Thr Arg Tyr Leu		
705	710	715
		720
Thr Arg Pro Leu		

<210> 5  
 <211> 588  
 <212> PRT  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/Note =  
synthetic construct

<400> 5

Thr	Ala	Pro	Thr	Gly	Lys	Arg	Ile	Asp	Asp	His	Phe	Pro	Lys	Arg	Lys
1				5					10					15	
Lys	Ala	Arg	Thr	Glu	Glu	Asp	Ser	Lys	Pro	Ser	Thr	Ser	Ser	Asp	Ala
			20					25					30		
Glu	Ala	Gly	Pro	Ser	Gly	Ser	Gln	Gln	Leu	Gln	Ile	Pro	Ala	Gln	Pro
		35					40					45			
Ala	Ser	Ser	Leu	Gly	Ala	Asp	Thr	Met	Ser	Ala	Gly	Gly	Gly	Gly	Pro
	50					55					60				
Leu	Gly	Asp	Asn	Asn	Gln	Gly	Ala	Asp	Gly	Val	Gly	Asn	Ala	Ser	Gly
65					70					75				80	
Asp	Trp	His	Cys	Asp	Ser	Thr	Trp	Met	Gly	Asp	Arg	Val	Val	Thr	Lys
			85						90					95	
Ser	Thr	Arg	Thr	Trp	Val	Leu	Pro	Ser	Tyr	Asn	Asn	His	Gln	Tyr	Arg
			100					105					110		
Glu	Ile	Lys	Ser	Gly	Ser	Val	Asp	Gly	Ser	Asn	Ala	Asn	Ala	Tyr	Phe
		115					120					125			
Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr	Phe	Asp	Phe	Asn	Arg	Phe	His	Ser
	130					135					140				
His	Trp	Ser	Pro	Arg	Asp	Trp	Gln	Arg	Leu	Ile	Asn	Asn	Tyr	Trp	Gly
145					150					155					160
Phe	Arg	Pro	Arg	Ser	Leu	Arg	Val	Lys	Ile	Phe	Asn	Ile	Gln	Val	Lys
				165					170					175	
Glu	Val	Thr	Val	Gln	Asp	Ser	Thr	Thr	Thr	Ile	Ala	Asn	Asn	Leu	Thr
			180					185					190		
Ser	Thr	Val	Gln	Val	Phe	Thr	Asp	Asp	Asp	Tyr	Gln	Leu	Pro	Tyr	Val
		195					200					205			
Val	Gly	Asn	Gly	Thr	Glu	Gly	Cys	Leu	Pro	Ala	Phe	Pro	Pro	Gln	Val
	210					215					220				
Phe	Thr	Leu	Pro	Gln	Tyr	Gly	Tyr	Ala	Thr	Leu	Asn	Arg	Asp	Asn	Thr
225					230					235					240
Glu	Asn	Pro	Thr	Glu	Arg	Ser	Ser	Phe	Phe	Cys	Leu	Glu	Tyr	Phe	Pro
				245					250					255	
Ser	Lys	Met	Leu	Arg	Thr	Gly	Asn	Asn	Phe	Glu	Phe	Thr	Tyr	Asn	Phe
			260					265					270		
Glu	Glu	Val	Pro	Phe	His	Ser	Ser	Phe	Ala	Pro	Ser	Gln	Asn	Leu	Phe
		275					280					285			
Lys	Leu	Ala	Asn	Pro	Leu	Val	Asp	Gln	Tyr	Leu	Tyr	Arg	Phe	Val	Ser
	290					295					300				
Thr	Asn	Asn	Thr	Gly	Gly	Val	Gln	Phe	Asn	Lys	Asn	Leu	Ala	Gly	Arg
305					310					315					320
Tyr	Ala	Asn	Thr	Tyr	Lys	Asn	Trp	Phe	Pro	Gly	Pro	Met	Gly	Arg	Thr
				325					330					335	
Gln	Gly	Trp	Asn	Leu	Gly	Ser	Gly	Val	Asn	Arg	Ala	Ser	Val	Ser	Ala
			340					345					350		
Phe	Ala	Thr	Thr	Asn	Arg	Met	Glu	Leu	Glu	Gly	Ala	Ser	Tyr	Gln	Val
		355					360					365			
Pro	Pro	Gln	Pro	Asn	Gly	Met	Thr	Asn	Asn	Leu	Gln	Gly	Ser	Asn	Thr
		370				375					380				
Tyr	Ala	Leu	Glu	Asn	Thr	Met	Ile	Phe	Asn	Ser	Gln	Pro	Ala	Asn	Pro
385					390					395					400
Gly	Thr	Thr	Ala	Thr	Tyr	Leu	Glu	Gly	Asn	Met	Leu	Ile	Thr	Ser	Glu
			405						410					415	
Ser	Glu	Thr	Gln	Pro	Val	Asn	Arg	Val	Ala	Tyr	Asn	Val	Gly	Gly	Gln
			420					425					430		
Met	Ala	Thr	Asn	Asn	Gln	Ser	Ser	Thr	Thr	Ala	Pro	Ala	Thr	Gly	Thr
		435				440						445			
Tyr	Asn	Leu	Gln	Glu	Ile	Val	Pro	Gly	Ser	Val	Trp	Met	Glu	Arg	Asp

```

      450                      455                      460
Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro Glu Thr Gly Ala
465                      470                      475                      480
His Phe His Pro Ser Pro Ala Met Gly Gly Phe Gly Leu Lys His Pro
      485                      490                      495
Pro Pro Met Met Leu Ile Lys Asn Thr Pro Val Pro Gly Asn Ile Thr
      500                      505                      510
Ser Phe Ser Asp Val Pro Val Ser Ser Phe Ile Thr Gln Tyr Ser Thr
      515                      520                      525
Gly Gln Val Thr Val Glu Met Glu Trp Glu Leu Lys Lys Glu Asn Ser
      530                      535                      540
Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Asn Asn Tyr Asn Asp Pro
545                      550                      555                      560
Gln Phe Val Asp Phe Ala Pro Asp Ser Thr Gly Glu Tyr Arg Thr Thr
      565                      570                      575
Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
      580                      585

```

```

<210> 6
<211> 532
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence:/Note =
        synthetic construct

```

```

      <400> 6
Met Ser Ala Gly Gly Gly Gly Pro Leu Gly Asp Asn Asn Gln Gly Ala
 1                      5                      10                      15
Asp Gly Val Gly Asn Ala Ser Gly Asp Trp His Cys Asp Ser Thr Trp
      20                      25                      30
Met Gly Asp Arg Val Val Thr Lys Ser Thr Arg Thr Trp Val Leu Pro
      35                      40                      45

Ser Tyr Asn Asn His Gln Tyr Arg Glu Ile Lys Ser Gly Ser Val Asp
 50                      55                      60
Gly Ser Asn Ala Asn Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr
65                      70                      75                      80
Phe Asp Phe Asn Arg Phe His Ser His Trp Ser Pro Arg Asp Trp Gln
      85                      90                      95
Arg Leu Ile Asn Asn Tyr Trp Gly Phe Arg Pro Arg Ser Leu Arg Val
      100                      105                      110
Lys Ile Phe Asn Ile Gln Val Lys Glu Val Thr Val Gln Asp Ser Thr
      115                      120                      125
Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp
      130                      135                      140
Asp Asp Tyr Gln Leu Pro Tyr Val Val Gly Asn Gly Thr Glu Gly Cys
145                      150                      155                      160
Leu Pro Ala Phe Pro Pro Gln Val Phe Thr Leu Pro Gln Tyr Gly Tyr
      165                      170                      175
Ala Thr Leu Asn Arg Asp Asn Thr Glu Asn Pro Thr Glu Arg Ser Ser
      180                      185                      190
Phe Phe Cys Leu Glu Tyr Phe Pro Ser Lys Met Leu Arg Thr Gly Asn
195                      200                      205
Asn Phe Glu Phe Thr Tyr Asn Phe Glu Glu Val Pro Phe His Ser Ser
      210                      215                      220
Phe Ala Pro Ser Gln Asn Leu Phe Lys Leu Ala Asn Pro Leu Val Asp
225                      230                      235                      240
Gln Tyr Leu Tyr Arg Phe Val Ser Thr Asn Asn Thr Gly Gly Val Gln
      245                      250                      255
Phe Asn Lys Asn Leu Ala Gly Arg Tyr Ala Asn Thr Tyr Lys Asn Trp
260                      265                      270

```

Phe Pro Gly Pro Met Gly Arg Thr Gln Gly Trp Asn Leu Gly Ser Gly  
 275 280 285  
 Val Asn Arg Ala Ser Val Ser Ala Phe Ala Thr Thr Asn Arg Met Glu  
 290 295 300  
 Leu Glu Gly Ala Ser Tyr Gln Val Pro Pro Gln Pro Asn Gly Met Thr  
 305 310 315 320  
 Asn Asn Leu Gln Gly Ser Asn Thr Tyr Ala Leu Glu Asn Thr Met Ile  
 325 330 335  
 Phe Asn Ser Gln Pro Ala Asn Pro Gly Thr Thr Ala Thr Tyr Leu Glu  
 340 345 350  
 Gly Asn Met Leu Ile Thr Ser Glu Ser Glu Thr Gln Pro Val Asn Arg  
 355 360 365  
 Val Ala Tyr Asn Val Gly Gly Gln Met Ala Thr Asn Asn Gln Ser Ser  
 370 375 380  
 Thr Thr Ala Pro Ala Thr Gly Thr Tyr Asn Leu Gln Glu Ile Val Pro  
 385 390 395 400  
 Gly Ser Val Trp Met Glu Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp  
 405 410 415  
 Ala Lys Ile Pro Glu Thr Gly Ala His Phe His Pro Ser Pro Ala Met  
 420 425 430  
 Gly Gly Phe Gly Leu Lys His Pro Pro Pro Met Met Leu Ile Lys Asn  
 435 440 445  
 Thr Pro Val Pro Gly Asn Ile Thr Ser Phe Ser Asp Val Pro Val Ser  
 450 455 460  
  
 Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Thr Val Glu Met Glu  
 465 470 475 480  
 Trp Glu Leu Lys Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln  
 485 490 495  
 Tyr Thr Asn Asn Tyr Asn Asp Pro Gln Phe Val Asp Phe Ala Pro Asp  
 500 505 510  
 Ser Thr Gly Glu Tyr Arg Thr Thr Arg Pro Ile Gly Thr Arg Tyr Leu  
 515 520 525  
 Thr Arg Pro Leu  
 530

&lt;210&gt; 7

&lt;211&gt; 2307

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

 <223> Description of Artificial Sequence: /Note =  
 synthetic construct

&lt;400&gt; 7

```

aggctctcat ttgttcccga gacgcctcgc agttcagacg tgactgttga tcccgcctcct      60
ctgcgaccgc tcaattggaa ttcaagtaaa taaagcgagt agtcatgtct tttgttgatc      120
accctccaga ttggttgga gaagttggtg aaggtcttcg cgagtttttg ggccttgaag      180
cgggccacc gaaacaaaaa cccaatcagc agcatcaaga tcaagcccg tggctttgtgc      240
tgcttggtta taactatctc ggacccggaa acggtctcga tcgaggagag cctgtcaaca      300
gggcagacga ggtcgcgcga gagcacgaca tctcgtacaa cgagcagctt gaggcgggag      360
acaacccta cctcaagtac aaccacgcgg acgccagatt tcaggagaag ctgcgccgacg      420
acacatcctt cgggggaaac ctcggaagg cagtctttca ggccaagaaa agggttctcg      480
aaccttttg cctggttgaa gaggttgcta agacggcccc taccgaaaag cggatagacg      540
accactttcc aaaaagaaag aaggctcggg ccgaagagga ctccaagcct tccacctcgt      600
cagacgccga agctggaccc agcggatccc agcagctgca aatcccagcc caaccagcct      660
caagtttggg agctgatata atgtctgcgg gaggtggcgg cccattgggc gacaataacc      720
aaggtgccga tggagtgggc aatgcctcgg gagattggca ttgcgattcc acgtggattg      780
gggacagagt cgtcaccaag tccaccgaa cctgggtgct gccagctac aacaaccacc      840
agtaccgaga gatcaaaagc ggctccgctg acggaagcaa cgccaacgcc tactttggat      900
acagcacccc ctgggggtac tttgacttta accgcttcca cagccactgg agccccgag      960
actggcaaag actcatcaac aactactggg gcttcagacc ccggtcctc agagtcaaaa     1020

```

tcttcaacat	tcaagtcaaa	gaggtcacgg	tgcaggactc	caccaccacc	atcgccaaca	1080
acctcacctc	caccgtccaa	gtgttttacg	acgacgacta	ccagctgccc	tacgtcgctg	1140
gcaacgggac	cgagggatgc	ctgccggcct	tccctccgca	ggtctttacg	ctgccgcagt	1200
acggttacgc	gacgctgaac	cgcgacaaca	cagaaaatcc	caccgagagg	agcagcttct	1260
tctgcctaga	gtactttccc	agcaagatgc	tgagaacggg	caacaacttt	gagtttacct	1320
acaactttga	ggaggtgccc	ttccactcca	gcttcgctcc	cagtcagaac	ctgttcaagc	1380
tggccaaccc	gctggtggac	cagtacttgt	accgcttcgt	gagcacaat	aacactggcg	1440
gagtcacagt	caacaagaac	ctggccggga	gatacgccaa	cacctacaaa	aactggttcc	1500
cggggcccat	gggcccgaac	cagggctgga	acctgggctc	cggggccaac	cgcgccagt	1560
tcagcgctt	cggcacgacc	aataggatgg	agctcgaggg	cgcgagttac	caggtgcccc	1620
cgcagccgaa	cggcatgacc	aacaacctcc	agggcagcaa	cacctatgcc	ctggagaaca	1680
ctatgatctt	caacagccag	ccggcgaacc	cgggcaccac	cgccacgtac	ctcgagggca	1740
acatgctcat	caccagcgag	agcgagacgc	agccggtgaa	ccgcgtggcg	tacaacgtcg	1800
gcgggcagat	ggccaccaac	aaccagagct	ccaccactgc	ccccgcgacc	ggcacgtaca	1860
acctccagga	aatcgtgccc	ggcagcgtgt	ggatggagag	ggacgtgtac	ctccaaggac	1920
ccatctgggc	caagatccca	gagacggggg	cgcactttca	cccctctccg	gccatggcg	1980
gattcggact	caaacacca	ccgccatga	tgctcatcaa	gaacacgcct	gtgcccgaa	2040
atatcaccag	cttctcggac	gtgcccgtca	gcagcttcat	caccagtag	agcaccgggc	2100
aggtcaccgt	ggagatggag	tgggagctca	agaaggaaaa	ctccaaggag	tggaaaccag	2160
agatccagta	cacaaacaac	tacaacgacc	cccagtttgt	ggactttgcc	ccggacagca	2220
ccggggaata	cagaaccacc	agacctatcg	gaacccgata	ccttaccgca	cccctttaac	2280
ccattcatgt	cgcataccct	caataaa				2307

&lt;210&gt; 8

&lt;211&gt; 2264

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: /Note =  
synthetic construct

&lt;400&gt; 8

aggctctcat	ttgttcccga	gacgcctcgc	agttcagacg	tgactgttga	tcccgtcct	60
ctgcgaccgc	tcaattggaa	ttcaagattg	gttgggaagaa	gttgggtgaag	gtcttcgca	120
gtttttgggc	cttgaagcgg	gcccaccgaa	acaaaaacc	aatcagcagc	atcaagatca	180
agcccgtggt	cttgtgctgc	ctggttataa	ctatctcgga	cccggaaacg	gtctcgatcg	240
aggagagcct	gtcaacaggg	cagacagagg	cgcgcgagag	cacgacatct	cgtacaacga	300
gcagcttgag	gcgggagaca	acccctacct	caagtacaac	cacgcggacg	ccgagtttca	360
ggagaagctc	gccgacgaca	catccttcgg	gggaaacctc	ggaaaggcag	tctttcaggc	420
caagaaaagg	gttctcgaac	cttttggcct	ggttgaagag	ggtgctaaga	cggccctac	480
cggaaagcgg	atagacgacc	actttccaaa	aagaaagaag	gctcggaccg	aagaggactc	540
caagccttcc	acctcgtcag	acgcccgaagc	tggaccagc	ggatcccagc	agctgcaaat	600
cccagcccaa	ccagcctcaa	gtttgggagc	tgatataatg	tctgcgggag	gtggcgcccc	660
attgggcgac	aataaccaag	gtgccgatgg	agtgggcaat	gcctcgggag	attggcattg	720
cgattccacg	tggatggggg	acagagtcgt	caccaagtcc	accggaacct	gggtgctgcc	780
cagctacaac	aaccaccagt	accgagagat	caaaagcggc	tccgtcgacg	gaagcaacgc	840
caacgcctac	tttgataaca	gcaccccttg	gggttacttt	gactttaacc	gcttccacag	900
ccactggagc	ccccgagact	ggcaaagact	catcaacaac	tactggggct	tcagacccc	960
gtccctcaga	gtcaaaatct	tcaacattca	agtcaaagag	gtcacgggtc	aggactccac	1020
caccaccatc	gccaacaacc	tcacctccac	cgtccaagtg	tttacggacg	acgactacca	1080
gctgccctac	gtcgtcggca	acgggaccga	gggatgcctg	ccggccttcc	ctccgcaggt	1140
ctttacgctg	ccgcagtagc	gttacgcgac	gctgaaccgc	gacaacacag	aaaatcccac	1200
cgagaggagc	agcttcttct	gcctagagta	ctttccagc	aagatgctga	gaacgggcaa	1260
caacttttag	tttacctaca	actttgagga	ggtgcccttc	cactccagct	tcgctccag	1320
tcagaacctg	ttcaagctgg	ccaaccctgt	ggtggaccag	tacttgtacc	gcttcgtgag	1380
cacaaataac	actggcggag	tccagttcaa	caagaacctg	gccgggagat	acgccaacac	1440
ctacaaaaac	tggttcccgg	ggcccatggg	ccgaacctag	ggctggaacc	tgggctccgg	1500
ggtcaaccgc	gccagtgtca	gcgccttcgc	cacgaccaat	aggatggagc	tcgagggcgc	1560
gagttaccag	gtgccccgc	agccgaacgg	catgaccaac	aacctccagg	gcagcaaac	1620
ctatgccctg	gagaacacta	tgatcttcaa	cagccagccg	gcgaacctgg	gcaccaccgc	1680
cacgtacctc	gagggcaaca	tgctcatcac	cagcgagagc	gagacgcagc	cggtagaccg	1740
cgtggcgtac	aacgtcggcg	ggcagatggc	caccaacaac	cagagctoca	ccactgcccc	1800

cgcgaccggc	acgtacaacc	tccaggaaat	cgtgcccggc	agcgtgtgga	tggagagggga	1860
cgtgtacctc	caaggaccca	tctggggccaa	gatcccagag	acggggggcgc	actttcaccc	1920
ctctccggcc	atggggcggat	tccgactcaa	acacccaccg	cccatgatgc	tcatcaagaa	1980
cacgcctgtg	cccggaaata	tcaccagctt	ctcggacgtg	cccgtcagca	gcttcatcac	2040
ccagtacagc	accggggcagg	tcaccgtgga	gatggagtgg	gagctcaaga	aggaaaactc	2100
caagaggtgg	aaccagaga	tccagtacac	aaacaactac	aacgaccccc	agtttgtgga	2160
ctttgccccg	gacagcaccg	gggaatacag	aaccaccaga	cctatcgga	cccgtacac	2220
taccgaccc	ctttaaccca	ttcatgtcgc	ataccctcaa	taaa		2264

<210> 9  
 <211> 2264  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 synthetic construct

<400> 9						
aggctctcat	ttgttcccga	gacgcctcgc	agttcagacg	tgactgttga	tcccgtcct	60
ctgcgaccgc	tcaattggaa	ttcaagattg	gttgggaagaa	gttgggtgaag	gtcttcgcga	120
gtttttgggc	cttgaagcgg	gcccaccgaa	acaaaaaccc	aatcagcagc	atcaagatca	180
agcccgtggt	cttgtgtctg	ctgggtataa	ctatctcgga	cccggaaacg	gtctcgatcg	240
aggagagcct	gtcaacaggg	cagacgaggt	cgcgcgagag	cacgacatct	cgtacaacga	300
gcagcttgag	gcgggagaca	acccctacct	caagtacaac	cacgcgagcg	ccgagtttca	360
ggagaagctc	gccgacgaca	catccttcgg	gggaaacctc	ggaaaggcag	tctttcagge	420
caagaaaagg	gttctcgaac	cttttggcct	ggttgaagag	ggtgctaaga	cggcccctac	480
cggaaagcgg	atagacgacc	actttccaaa	aagaaagaag	gctcggaccg	aagaggactc	540
caagccttcc	acctcgtcag	acgccgaagc	tggaccacgc	ggatcccagc	agctgcaaat	600
cccagcccaa	ccagcctcaa	gttttgggagc	tgatacaatg	tctgcgggag	gtggcgcccc	660
attgggcgac	aataaccaag	gtgccgatgg	agtgggcaat	gcctcgggag	attggcattg	720
cgattccacg	tggatggggg	acagagtcgt	caccaagtc	acccgaacct	gggtgtctgc	780
cagctacaac	aaccaccagt	acogagagat	caaaagcggc	tccgtcgacg	gaagcaacgc	840
caacgcctac	tttgatata	gcacccctg	gggtacttt	gaatttaacc	gcttccacag	900
ccactggagc	ccccgagact	ggcaaagact	catcaacaac	tactggggct	tcagaccccc	960
gtccctcaga	gtcaaaatct	tcaacattca	agtcaaagag	gtcacgggtg	aggactccac	1020
caccaccatc	gccaacaacc	tcacctccac	cgtccaagt	tttacggacg	acgactacca	1080
gctgccctac	gtcgtcgga	acgggaccga	gggatgccctg	ccggccttcc	ctccgcaggt	1140
ctttacgctg	ccgcagtacg	gttacgcgac	gctgaaccgc	gacaacacag	aaaatcccac	1200
cgagaggagc	agcttcttct	gcctagagta	ctttccagc	aagatgctga	gaacgggcaa	1260
caactttgag	tttacctaca	actttgagga	ggtgcccttc	cactccagct	tcgctcccag	1320
tcagaacctg	ttcaagctgg	ccaacccgct	ggtggaccag	tacttgtacc	gcttcgtgag	1380
cacaaataac	actggcggag	tccagttcaa	caagaacctg	gccgggagat	acgcacaac	1440
ctacaaaaac	tggttccggg	ggcccatggg	ccgaacccag	ggctggaacc	tgggtctcgg	1500
ggtcaaccgc	gccagtgtca	gcgccttcgc	cacgaccaat	aggatggagc	tcgagggcgc	1560
gagttaccag	gtgccccgcg	agccgaacgg	catgaccaac	aacctccagg	gcagcaacac	1620
ctatgcctg	gagaacacta	tgatcttcaa	cagccagccg	gcgaacccgg	gcaccaccgc	1680
cacgtacctc	gagggcaaca	tgctcatcac	cagcgagagc	gagacgcagc	cgggtgaaccg	1740
cgtggcgtag	aacgtcggcg	ggcagatggc	caccaacaac	cagagctcca	ccactgcccc	1800
cgcgaccggc	acgtacaacc	tccaggaaat	cgtgcccggc	agcgtgtgga	tggagagggga	1860
cgtgtacctc	caaggaccca	tctggggccaa	gatcccagag	acggggggcgc	actttcaccc	1920
ctctccggcc	atggggcggat	tccgactcaa	acacccaccg	cccatgatgc	tcatcaagaa	1980
cacgcctgtg	cccggaaata	tcaccagctt	ctcggacgtg	cccgtcagca	gcttcatcac	2040
ccagtacagc	accgggcagg	tcaccgtgga	gatggagtgg	gagctcaaga	aggaaaactc	2100
caagaggtgg	aaccagaga	tccagtacac	aaacaactac	aacgaccccc	agtttgtgga	2160
ctttgccccg	gacagcaccg	gggaatacag	aaccaccaga	cctatcgga	cccgtacac	2220
taccgaccc	ctttaaccca	ttcatgtcgc	ataccctcaa	taaa		2264

<210> 10  
 <211> 1292  
 <212> DNA  
 <213> Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
synthetic construct

&lt;400&gt; 10

agcgcgcaaacg	gctcgtcgcg	cagtttcttg	cagaatcttc	gcagcgctcg	caggaggcgg	60
cttcgcagcg	tgagttctcg	gctgaccccg	tcatacaaaag	caagacttcc	cagaaataca	120
tggcgctcgt	caactggctc	gtggagcacg	gcatacattc	cgagaagcag	tgatccagg	180
aaaatcagga	gagctacctc	tccttcaact	ccaccggcaa	ctctcggagc	cagatcaagg	240
ccgcgctcga	caacgcgacc	aaaattatga	gtctgacaaa	aagcgcggtg	gactacctcg	300
tggggagctc	cgttcccgag	gacatttcaa	aaaacagaat	ctggcacaatt	tttgagatga	360
atggctacga	cccggcctac	gcgggatcca	tcctctacgg	ctggtgtcag	cgctccttca	420
acaagaggaa	caccgtctgg	ctctacggac	ccgccacgac	cggcaagacc	aacatcgagg	480
aggccatcgc	ccacactgtg	cccttttacg	gctgcgtgaa	ctggaccaat	gaaaactttc	540
cctttaatga	ctgtgtggac	aaaatgctca	tttgggtgga	ggagggaag	atgaccaaca	600
agggtggtga	atccgccaaag	gccatcctgg	ggggctcaaa	ggtgcgggtc	gatcagaaat	660
gtaaatcctc	tgttcaaat	gattctaccc	ctgtcattgt	aacttccaat	acaaacatgt	720
gtgtggtggt	ggatgggaat	tccacgacct	ttgaacacca	gcagccgctg	gaggaccgca	780
tgttcaaat	tgaactgact	aagcggtccc	cgccagattt	tggcaagatt	actaagcagg	840
aagtcaagga	cttttttgc	tgggcaagg	tcaatcaggt	gccggtgact	cacgagttta	900
aagttcccg	ggaattggcg	ggaactaaag	gggaggagaa	atctctaaaa	cgcccactgg	960
gtgacgtcac	caatactagc	tataaaagtc	tggagaagcg	ggccaggctc	tcatttggtc	1020
ccgagacgcc	tcgcagttca	gacgtgactg	ttgatccgcg	tcctctgcga	ccgctcaatt	1080
ggaattcaag	gtatgattgc	aaatgtgact	atcatgctca	atttgacaa	atttctaaca	1140
aatgtgatga	atgtgaatat	ttgaatcggg	gcaaaaatgg	atgtatctgt	cacaatgtaa	1200
ctcactgtca	aatttgtcat	gggattcccc	cctgggaaaa	ggaaaacttg	tcagattttg	1260
gggattttga	cgatgccaat	aaagaacagt	aa			1292

&lt;210&gt; 11

&lt;211&gt; 1870

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
synthetic construct

&lt;400&gt; 11

attcttttgc	ctggactgct	agaggaccct	cgctgccatg	gctaccttct	atgaagtcac	60
tgttcgcgtc	ccatttgacg	tggaggaaca	tctgcctgga	atttctgaca	gctttgtgga	120
ctgggtaact	gggtcaaat	gggagctgcc	tccagagtc	gattttaa	tgactctggt	180
tgaacagcct	cagttgacgg	tggctgatag	aattcgccgc	gtgttcctgt	acgagtggaa	240
caaattttcc	aagcaggagt	ccaaattctt	tgtgcagttt	gaaaagggat	ctgaatat	300
tcactctgac	acgcttggg	agacctccgg	catctcttcc	atggctcctg	gccgtacgt	360
gagtcagatt	cgcgcccagc	tgggtgaaagt	gggtctccag	ggaattgaac	cccagatcaa	420
cgactgggtc	gccatcacca	aggtaaagaa	gggaggagcc	aataagggtg	tggattctgg	480
gtatattccc	gcctacctgc	tgccgaaggt	ccaaccggag	cttcagtg	cgtggacaaa	540
cctggacgag	tataaattgg	ccgccctgaa	tctggaggag	cgaaaacggc	tcgtcgcgca	600
gtttctggca	gaatcctcgc	agcgctcgca	ggaggcggtc	tcgcagcggtg	agttctcggc	660
tgacccggtc	atcaaaaagca	agacttccca	gaaatacatg	gcgctcgtca	actggctcgt	720
ggagcacggc	atcacttccg	agaagcagtg	gatccaggaa	aatcaggaga	gctacctctc	780
cttcaactcc	accggcaact	ctcggagcca	gatcaaggcc	gcgctcgaca	acgcgaccaa	840
aattatgagt	ctgacaaaaa	gcgcgggtgga	ctacctcgtg	gggagctccg	ttcccagga	900
catttcaaaa	aacagaatct	ggcaaat	tgagatgaat	ggctacgacc	cggcctacgc	960
gggatccatc	ctctacggct	ggtgtcagcg	ctccttcaac	aagaggaaca	ccgtctggct	1020
ctacggaccc	gccacgaccg	gcaagaccaa	catcgcgag	gccatcgccc	acactgtgcc	1080
cttttacggc	tgcgtgaact	ggaccaatga	aaactttccc	tttaattgact	gtgtggacaa	1140
aatgtcatt	tgggtgggag	agggaaagat	gaccaacaag	gtggttgaat	ccgccaaagg	1200
catcctgggg	ggctcaaagg	tgcgggtcga	tcagaaatgt	aaatcctctg	ttcaaattga	1260
ttctaccct	gtcattgtaa	cttccaatac	aaacatgtgt	gtggtggtgg	atgggaattc	1320
cacgaccttt	gaacaccagc	agccgctgga	ggaccgcatg	ttcaaat	aactgactaa	1380
gcggctcccc	ccagattttg	gcaagattac	taagcaggaa	gtcaaggact	tttttgcttg	1440
ggcaagggtc	aatcagggtc	cggtgactca	cgagtttaaa	gttcccaggg	aattggcggg	1500



13

```

aactaaaggg ggggagaaat ctctaaaacg cccactgggt gacgtcacca atactagcta 1560
taaaagtctg gagaagcggg ccaggctctc atttggtccc gagacgcctc gcagttcaga 1620
cgtgactggt gatccccgtc ctctgcgacc gctcaattgg aattcaaggt atgattgcaa 1680
atgtgactat catgctcaat ttgacaacat ttctaacaaa tgtgatgaat gtgaatattt 1740
gaatcggggc aaaaatggat gtatctgtca caatgtaact cactgtcaaa ttgtcatgg 1800
gattccccc tgggaaaagg aaaacttgtc agattttggg gattttgacg atgccaataa 1860
agaacagtaa 1870

```

<210> 12  
 <211> 330  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 synthetic construct

<400> 12

Met	Ala	Leu	Val	Asn	Trp	Leu	Val	Glu	His	Gly	Ile	Thr	Ser	Glu	Lys
1				5					10					15	
Gln	Trp	Ile	Gln	Asn	Gln	Glu	Ser	Tyr	Leu	Ser	Phe	Asn	Ser	Thr	
			20					25					30		
Gly	Asn	Ser	Arg	Ser	Gln	Ile	Lys	Ala	Ala	Leu	Asp	Asn	Ala	Thr	Lys
			35				40						45		
Ile	Met	Ser	Leu	Thr	Lys	Ser	Ala	Val	Asp	Tyr	Leu	Val	Gly	Ser	Ser
	50					55					60				
Val	Pro	Glu	Asp	Ile	Ser	Lys	Asn	Arg	Ile	Trp	Gln	Ile	Phe	Glu	Met
65						70				75					80
Asn	Gly	Tyr	Asp	Pro	Ala	Tyr	Ala	Gly	Ser	Ile	Leu	Tyr	Gly	Trp	Cys
				85					90					95	
Gln	Arg	Ser	Phe	Asn	Lys	Arg	Asn	Thr	Val	Trp	Leu	Tyr	Gly	Pro	Ala
			100					105					110		
Thr	Thr	Gly	Lys	Thr	Asn	Ile	Ala	Glu	Ala	Ile	Ala	His	Thr	Val	Pro
		115					120					125			
Phe	Tyr	Gly	Cys	Val	Asn	Trp	Thr	Asn	Glu	Asn	Phe	Pro	Phe	Asn	Asp
		130				135					140				
Cys	Val	Asp	Lys	Met	Leu	Ile	Trp	Trp	Glu	Glu	Gly	Lys	Met	Thr	Asn
145					150					155					160
Lys	Val	Val	Glu	Ser	Ala	Lys	Ala	Ile	Leu	Gly	Gly	Ser	Lys	Val	Arg
				165					170					175	
Val	Asp	Gln	Lys	Cys	Lys	Ser	Ser	Val	Gln	Ile	Asp	Ser	Thr	Pro	Val
			180					185					190		
Ile	Val	Thr	Ser	Asn	Thr	Asn	Met	Cys	Val	Val	Val	Asp	Gly	Asn	Ser
		195				200						205			
Thr	Thr	Phe	Glu	His	Gln	Gln	Pro	Leu	Glu	Asp	Arg	Met	Phe	Lys	Phe
		210				215					220				
Glu	Leu	Thr	Lys	Arg	Leu	Pro	Pro	Asp	Phe	Gly	Lys	Ile	Thr	Lys	Gln
225					230					235					240
Glu	Val	Lys	Asp	Phe	Phe	Ala	Trp	Ala	Lys	Val	Asn	Gln	Val	Pro	Val
				245					250					255	
Thr	His	Glu	Phe	Lys	Val	Pro	Arg	Glu	Leu	Ala	Gly	Thr	Lys	Gly	Ala
			260					265					270		
Glu	Lys	Ser	Leu	Lys	Arg	Pro	Leu	Gly	Asp	Val	Thr	Asn	Thr	Ser	Tyr
		275				280						285			
Lys	Ser	Leu	Glu	Lys	Arg	Ala	Arg	Leu	Ser	Phe	Val	Pro	Glu	Thr	Pro
		290				295					300				
Arg	Ser	Ser	Asp	Val	Thr	Val	Asp	Pro	Ala	Pro	Leu	Arg	Pro	Leu	Asn
305					310					315					320
Trp	Asn	Ser	Arg	Leu	Val	Gly	Arg	Ser	Trp						
				325					330						

<210> 13  
 <211> 1115

14

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
synthetic construct

&lt;400&gt; 13

```

aggagcgcgca acggctcgct ggcgagtttc tggcagaatc ctgcgacgcg tcgcaggagg      60
cggttcgcga gcgtgagttc tcggctgacc cggatcatcaa aagcaagact tcccagaaat      120
acatggcgct cgtcaactgg ctcgtaggagc acggcatcac ttccgagaag cagtggatcc      180
aggaaaatca ggagagctac ctctccttca actccaccgg caactctcgg agccagatca      240
aggccgcgct cgacaacgcg accaaaatta tgagtctgac aaaaagcgcg gtggactacc      300
tcgtggggag ctccggttccc gaggacattt caaaaaacag aatctggcaa atttttgaga      360
tgaatggcta cgaccgggcc tacgcgggat ccatacctcta cggttggtgt cagcgctcct      420
tcaacaagag gaacaccgct tggctctacg gaccggccac gaccggcaag accaacatcg      480
cggaggccat cgcccacact gtgccctttt acggctgcbt gaactggacc aatgaaaact      540
ttccctttaa tgactgtgtg gacaaaatgc tcatttggtg ggaggaggga aagatgacca      600
acaaggtggt tgaatccgcc aaggccatcc tggggggctc aaaggtgcgg gtcgatcaga      660
aatgtaaatc ctctgttcaa attgattcta cccctgtcat tgaacttcc aatacaaaaca      720
tgtgtgtggt ggtggatggg aattccacga cctttgaaca ccagcagccg ctggaggacc      780
gcatgttcaa atttgaactg actaagcgcc tcccgccaga ttttgcaag attactaagc      840
aggaagtcaa ggactttttt gcttgggcaa aggtcaatca ggtgccggtg actcacgagt      900
ttaaagttcc caggaattg gcgggaacta aaggggcgga gaaatctcta aaacgcccac      960
tgggtgacgt caccaatact agctataaaa gtctggagaa gcggggccagg ctctcatttg     1020
ttcccgagac gcctcgcatg tcagacgtga ctgttgatcc cgctcctctg cgaccgctca     1080
attggaattc aagattggtt ggaagaagtt ggtga                                     1115

```

&lt;210&gt; 14

&lt;211&gt; 550

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
synthetic construct

&lt;400&gt; 14

```

Met Ala Thr Phe Tyr Glu Val Ile Val Arg Val Pro Phe Asp Val Glu
 1          5          10          15
Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Asp Trp Val Thr Gly
 20          25          30
Gln Ile Trp Glu Leu Pro Pro Glu Ser Asp Leu Asn Leu Thr Leu Val
 35          40          45
Glu Gln Pro Gln Leu Thr Val Ala Asp Arg Ile Arg Arg Val Phe Leu
 50          55          60
Tyr Glu Trp Asn Lys Phe Ser Lys Gln Glu Ser Lys Phe Phe Val Gln
 65          70          75          80
Phe Glu Lys Gly Ser Glu Tyr Phe His Leu His Thr Leu Val Glu Thr
 85          90          95
Ser Gly Ile Ser Ser Met Val Leu Gly Arg Tyr Val Ser Gln Ile Arg
100          105          110
Ala Gln Leu Val Lys Val Val Phe Gln Gly Ile Glu Pro Gln Ile Asn
115          120          125
Asp Trp Val Ala Ile Thr Lys Val Lys Lys Gly Gly Ala Asn Lys Val
130          135          140
Val Asp Ser Gly Tyr Ile Pro Ala Tyr Leu Leu Pro Lys Val Gln Pro
145          150          155          160
Glu Leu Gln Trp Ala Trp Thr Asn Leu Asp Glu Tyr Lys Leu Ala Ala
165          170          175
Leu Asn Leu Glu Glu Arg Lys Arg Leu Val Ala Gln Phe Leu Ala Glu
180          185          190
Ser Ser Gln Arg Ser Gln Glu Ala Ala Ser Gln Arg Glu Phe Ser Ala

```

15

195					200					205					
Asp	Pro	Val	Ile	Lys	Ser	Lys	Thr	Ser	Gln	Lys	Tyr	Met	Ala	Leu	Val
210						215					220				
Asn	Trp	Leu	Val	Glu	His	Gly	Ile	Thr	Ser	Glu	Lys	Gln	Trp	Ile	Gln
225					230					235					240
Glu	Asn	Gln	Glu	Ser	Tyr	Leu	Ser	Phe	Asn	Ser	Thr	Gly	Asn	Ser	Arg
				245					250					255	
Ser	Gln	Ile	Lys	Ala	Ala	Leu	Asp	Asn	Ala	Thr	Lys	Ile	Met	Ser	Leu
			260				265						270		
Thr	Lys	Ser	Ala	Val	Asp	Tyr	Leu	Val	Gly	Ser	Ser	Val	Pro	Glu	Asp
			275				280					285			
Ile	Ser	Lys	Asn	Arg	Ile	Trp	Gln	Ile	Phe	Glu	Met	Asn	Gly	Tyr	Asp
			290			295					300				
Pro	Ala	Tyr	Ala	Gly	Ser	Ile	Leu	Tyr	Gly	Trp	Cys	Gln	Arg	Ser	Phe
305					310					315					320
Asn	Lys	Arg	Asn	Thr	Val	Trp	Leu	Tyr	Gly	Pro	Ala	Thr	Thr	Gly	Lys
			325						330					335	
Thr	Asn	Ile	Ala	Glu	Ala	Ile	Ala	His	Thr	Val	Pro	Phe	Tyr	Gly	Cys
			340					345					350		
Val	Asn	Trp	Thr	Asn	Glu	Asn	Phe	Pro	Phe	Asn	Asp	Cys	Val	Asp	Lys
			355				360					365			
Met	Leu	Ile	Trp	Trp	Glu	Glu	Gly	Lys	Met	Thr	Asn	Lys	Val	Val	Glu
			370			375					380				
Ser	Ala	Lys	Ala	Ile	Leu	Gly	Gly	Ser	Lys	Val	Arg	Val	Asp	Gln	Lys
385					390					395					400
Cys	Lys	Ser	Ser	Val	Gln	Ile	Asp	Ser	Thr	Pro	Val	Ile	Val	Thr	Ser
			405						410					415	
Asn	Thr	Asn	Met	Cys	Val	Val	Val	Asp	Gly	Asn	Ser	Thr	Thr	Phe	Glu
			420					425					430		
His	Gln	Gln	Pro	Leu	Glu	Asp	Arg	Met	Phe	Lys	Phe	Glu	Leu	Thr	Lys
			435				440					445			
Arg	Leu	Pro	Pro	Asp	Phe	Gly	Lys	Ile	Thr	Lys	Gln	Glu	Val	Lys	Asp
			450			455					460				
Phe	Phe	Ala	Trp	Ala	Lys	Val	Asn	Gln	Val	Pro	Val	Thr	His	Glu	Phe
465					470					475					480
Lys	Val	Pro	Arg	Glu	Leu	Ala	Gly	Thr	Lys	Gly	Ala	Glu	Lys	Ser	Leu
			485					490					495		
Lys	Arg	Pro	Leu	Gly	Asp	Val	Thr	Asn	Thr	Ser	Tyr	Lys	Ser	Leu	Glu
			500				505					510			
Lys	Arg	Ala	Arg	Leu	Ser	Phe	Val	Pro	Glu	Thr	Pro	Arg	Ser	Ser	Asp
			515				520					525			
Val	Thr	Val	Asp	Pro	Ala	Pro	Leu	Arg	Pro	Leu	Asn	Trp	Asn	Ser	Arg
			530			535					540				
Leu	Val	Gly	Arg	Ser	Trp										
545					550										

<210> 15  
 <211> 1690  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 synthetic construct

<400> 15	
attctttgct ctggactgct agaggaccct cgctgccatg gctaccttct atgaagtcac	60
tggttcgcgct ccatttgacg tggaggaaca tctgcctgga atttctgaca gctttgtgga	120
ctgggtaact ggtcaaattt gggagctgcc tccagagtca gatttaaatt tgactctggt	180
tgaacagcct cagttgacgg tggctgatag aattcgccgc gtgttcctgt acgagtggaa	240
caaattttcc aagcaggagt ccaaattctt tgtgcagttt gaaaagggat ctgaatat	300
tcactctgac acgcttggtg agacctcgg catctcttcc atggtcctcg gccgctacgt	360

16

```

gagtcagatt cgcgcccagc tgggtgaaagt ggtcttccag ggaattgaac cccagatcaa 420
cgactgggtc gccatcacca aggttaaagaa gggcggagcc aataaggtgg tggattctgg 480
gtatattccc gcctacctgc tgccgaaggt ccaaccggag cttcagtggt cgtggacaaa 540
cctggacgag tataaattgg ccgccctgaa tctggaggag cgcaaacggc tcgtcgcgca 600
gtttctggca gaatcctcgc agcgcctcgca ggaggcggct tcgcagcgtg agttctcggc 660
tgacccggtc atcaaaagca agacttccca gaaatacatg gcgctcgtca actggctcgt 720
ggagcacggc atcacttccg agaagcagtg gatccaggaa aatcaggaga gctacctctc 780
cttcaactcc accggcaact ctccggagcca gatcaaggcc gcgctcgaca acgcgaccaa 840
aattatgagt ctgacaaaaa gcgcgggtgga ctacctcgtg gggagctccg ttcccaggga 900
catttcaaaa aacagaatct ggcaaatttt tgagatgaat ggctacgacc cggcctacgc 960
gggatccatc ctctacggct ggtgtcagcg ctcttcaac aagagggaaca ccgtctggct 1020
ctacggaccc gccacgaccg gcaagaccaa catcgcgagg gccatcgccc acactgtgcc 1080
cttttacggc tgcgtgaact ggaccaatga aaactttccc tttaatgact gtgtggacaa 1140
aatgctcatt tgggtgggag agggaaaagat gaccaacaag gtggttgaat ccgccaaagg 1200
catctggggg ggctcaaagg tgccgggtcga tcagaaatgt aaatcctctg ttcaaattga 1260
ttctaccctc gtcattgtaa ctccaatac aaacatgtgt gtggtgggtg atgggaattc 1320
cacgaccttt gaacaccagc agccgctgga ggaccgcatg ttcaaatttg aactgactaa 1380
gcggtctccg ccagattttg gcaagattac taagcaggaa gtcaaggact ttttgcttg 1440
ggcaaagggt aatcaggtgc cggtgactca cgagtttaaa gtcccagggt aattggcggg 1500
aactaaaggg gcggagaaat ctctaaaacg cccactgggt gacgtcacca atactagcta 1560
taaaagtctg gagaagcggg ccaggctctc atttgttccc gagacgcctc gcagttcaga 1620
cgtgactgtt gatccgcctc ctctgcgacc gctcaattgg aattcaagat tggttggaag 1680
aagttggtga

```

<210> 16  
 <211> 145  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 synthetic construct

```

<400> 16
ccatcaccaa ggtaaagaag ggcggagcca ataaggtggt ggattctggg tatattcccg 60
cctacctgct gccgaagggt caaccggagc ttcagtggtg gtggacaaaac ctggacgagt 120
ataaattggc cgccctgaat ctgga

```

<210> 17  
 <211> 174  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 synthetic construct

```

<400> 17
taagcaggaa gtcaaggact tttttgcttg ggcaaagggt aatcaggtgc cggtgactca 60
cgagtttaaa gttcccagggt aattggcggg aactaaaggg gcggagaaat ctctaaaacg 120
cccactgggt gacgtcacca atactagcta taaaagtctg gagaagcggg ccag 174

```

<210> 18  
 <211> 187  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 synthetic construct

```

<400> 18
cactctcaag caagggggtt ttgtaagcag tgatgtcata atgatgtaat gcttattgtc 60

```

17

acgcgatagt taatgattaa cagtcattgt atgtgtttta tccaatagga agaaagcgcg 120  
 cgtatgagtt ctgcgagac ttccggggta taaaagaccg agtgaacgag cccgcccga 180  
 ttctttg 187

<210> 19  
 <211> 168  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 synthetic construct

<400> 19  
 aaacctcctt gcttgagagt gtggcactct cccccctgtc gcgttcgctc gctcgtggc 60  
 tcgtttgggg ggggtggcagc tcaaagagct gccagacgac ggccctctgg ccgtcgcccc 120  
 cccaaacgag ccagcgagcg agcgaacgag acagggggga gaggcca 168

<210> 20  
 <211> 168  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 synthetic construct

<400> 20  
 aaacctcctt gcttgagagt gtggcactct cccccctgtc gcgttcgctc gctcgtggc 60  
 tcgtttgggg gggcgacggc cagagggccg tcgtctgccc gctctttgag ctgccacccc 120  
 cccaaacgag ccagcgagcg agcgaacgag acagggggga gaggcca 168

<210> 21  
 <211> 8  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 synthetic construct

<400> 21  
 cgggtgga 8

<210> 22  
 <211> 8  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 synthetic construct

<400> 22  
 cggttgag 8

<210> 23  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>

18

<223> Description of Artificial Sequence:/Note =  
synthetic construct

<400> 23  
caaaacctcc ttgcttgaga g

21